


Provided for non-commercial research and education use.

Not for reproduction, distribution or commercial use.




This article was published in an Sjournals journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the authors institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copied, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Text form) to their personal website or institutional repository. Authors requiring further information regarding Sjournals's archiving and manuscript policies encouraged to visit:

<http://www.sjournals.com>

© 2020 Sjournals Publishing Company



Contents lists available at Sjournals

## Agricultural Advances

Journal homepage: [www.sjournals.com](http://www.sjournals.com)

### Original article

## A comparative study in fish biodiversity in Goronyo reservoir and downstream river Rima in Sokoto State Northwestern Nigeria

Ibrahim Abdullahi Gusau<sup>a,\*</sup>, U. Muhammad<sup>b</sup>, B.D. Bilbis<sup>b</sup>

<sup>a</sup>Department of Fisheries and Aquaculture, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, Nigeria.

<sup>b</sup>National Biotechnology Development Agency (NABDA), Airport Road Lugbe, Abuja, Nigeria.

\*Corresponding author: [ibrahim.abdullahi@udusok.edu.ng](mailto:ibrahim.abdullahi@udusok.edu.ng)

### ARTICLE INFO

#### Article history,

Received 19 June 2020

Accepted 16 July 2020

Available online 23 July 2020

iThenticate screening 21 June 2020

English editing 15 July 2020

Quality control 21 July 2020

#### Keywords,

Reservoir

Rivers

Landing sites

Fish

Nigeria

### ABSTRACT

This study was conducted between April and October, 2016 in Goronyo Reservoir and downstream River Rima in Sokoto State with the main objective of evaluating and comparing the fish biodiversities of the two water bodies in relation to types of gears used. A multistage sampling technique was used to select water bodies. Three landing sites were purposely selected from downstream River Rima and five from Goronyo reservoir because of their high fish landings and five fishermen were randomly selected and their catches monitored. The catches were identified at species level, counted, measured and weighed to the nearest centimeter and gram respectively. The results showed the presence of 14 and 15 fish families in river Rima and Goronyo reservoir, represented by 40 and 35 fish species respectively. The members of Mormyridae and Bagridae families were the dominant fishes in terms of numbers accounting for 44.92%, 16.92% in River Rima and 26.06%, 21.44% in Goronyo reservoir. The most rarely occurring families were Clariidae and Cyprinidae in Goronyo reservoir accounting for 0.26% and 0.45% respectively. However, rarest species in Rima River was *Synodontis membranaceus* (0.11%). *Mormyrus rume* was the dominant species in both River Rima and Goronyo reservoir accounting for 23.64% and 15.31% by number. Simpson species richness showed Rima River with 40 and Goronyo reservoir with 35 fish species each exhibiting high (0.90) Simpson index of heterogeneity. Hook and line and gillnets were the most common fish gears used by the fishermen accounting for

---

29.87% and 30.37% respectively. Our study concluded that the two water bodies have fairly high fish diversities and recommended the reservoir should be enhanced by stocking it with *Lates niloticus* and *Labeo coubie* to serve as sport and commercial fishes in the reservoir.

© 2020 Sjournals. All rights reserved.

---

## 1. Introduction

Nigeria has a large spread of freshwater resources that exist in the form of lakes, reservoirs, canals and rivers. Freshwater is critical to human society and sustains all terrestrial and aquatic ecosystems (Millennium Ecosystem Assessment, 2005). The fish population of any water body that is capable of reproducing naturally in the wild and exploiting its fish diversity which varies from one water body and another (Ita, 1993). Biodiversity is a fundamental property of every living system that manifests at every level of hierarchy from molecular to ecosystem (Gupta and Gupta, 2008). Fish occurs in water bodies in different proportions and composition, in other words, Ichthyofauna composition of any water body, marine or fresh water differs in its fish species diversity and abundance (Ita, 1993; Joseph et al., 2015). Gupta and Gupta (2008) reported that fish exhibit the greatest diversity among the world vertebrates, with over 22,000-30,000 species and counting. The authors noted biodiversity is essential for balancing the ecosystem and facing varied problems to the environment.

Ita (1993), Dankishiya et al. (2013) and Josep et al. (2015) observed that a community of fish is said to have high species diversity if many equally or nearly equal abundant species are present. Conversely, if a community is composed of a very few species or if only a few species are abundant, then species diversity is low (Ibrahim et al., 2009). Gupta and Gupta (2008) further reported that high species diversity indicates a highly complex community, for a greater variety of species allows for a larger array of species interaction. Thus, population interaction involving energy transfer (food webs), predation, competition and niche apportionment are theoretically more complex and varied in a community of high species diversity (Joseph et al., 2015). Balogun (2005) and Joseph et al. (2015) reported that high species diversity correlates with community stability; the ability of community structure to be unaffected by disturbance of its components.

In a review of the Nigerian fisheries resources, Ita (1993) reported about 511 fish families from different aquatic habitats. Out of these fish families, about 22% of their fish species belong to brackish water and 44% is fresh water fish species inhabiting water of low salinity, (Ita, 1993). The author further reported that about 34% of the species are restricted to an exclusive economic zone (EEZ) covering about 220 nautical miles offshore.

The fish biodiversity in downstream River Rima in comparison to that of Goronyo reservoir was not adequately investigated. The information available is scanty and misleading, hence this paper aims at providing up to date information on fish species diversities in the two water bodies in relation to the fish gears used by the fishermen.

## 2. Materials and methods

### 2.1. Description of the study site/location

The study was conducted in River Rima and Goronyo reservoir all in Sokoto State, Northwestern Nigeria. The state lies between longitudes 5.14 0 E and 5.30 0 E and latitudes 12.96 0 N and 13.08 0 N. (Eniolorunda, 2012). The upper Rima was seasonal before it was dammed at Goronyo village, and flows during the rainy season overflowing its banks in August and September (Ita et al., 1982). The reservoir formed after damming the Rima river is known as Goronyo reservoir, it lies on coordinates 13° 31'50"N 05° 52'56"E (Google earth, 2016). The reservoir has a storage capacity of 976,000,000 cubic meters (Sembenelli Consulting, 2010). It has rivers Bunsuru, Maradi and Gangare as its main tributaries.

### 2.2. Data collection and sample size

A multistage sampling technique was used in this study. Two water bodies were purposely selected (downstream River Rima and Goronyo reservoir). Three landing sites from river Rima were selected (Usmanu

Danfodiyo University Bridge, Kwalkwalawa and Dundaye villages) and designated as stations A, B, and C. Additional five fish landing sites which comprised of Killaro, Rimawa, Keita, Katsira and Gidan yar fara also at Goronyo reservoir were also purposively selected because of their high fish landings and five fishermen were randomly selected from each station and landing site giving a total of 40 fishermen and their catches monitored. A total of 2,554 fish specimens were collected both at River Rima and Goronyo reservoir for data collection. The data was collected on monthly basis for a period of six months from April to October, 2016.

### 2.3. The fish identification and number

The number of individuals of the same fish family, genus and species was identified and counted according to the procedure described by Gusau et al. (2016).

### 2.4. The fish length/size

The length of the individual fish specimen was measured using a meter rule to the nearest mm expressed as total and standard lengths as described by Shinkafi et al. (2013).

### 2.5. The fish weight

The individual fish specimen of each species was weighed using a weighing balance to the nearest gram (g) as described by Bagenal and Tesch (1978).

### 2.6. Species density and relative abundance

The information obtained from the above procedures served as inputs based on the scale of the frequency of occurrence as described by Benech et al. (1983) for species dominance determination as follows:

Dominant = when a species occurred more than or equal to 10% ( $\geq 10\%$ ) in the population of fishes caught by the fishermen.

Sub-dominant = when species occurred from one percent to nine percent (1-9%) in the population of fishes caught by the fishermen.

Occasional = when a species occurred less than one or equal to one percent ( $\leq 1\%$ ) in the population of fishes caught by the fishermen.

Rare = when it was caught only once (1) in the population of fishes caught by the fishermen.

Extinct = when any species that was in the reservoir or river before but was not caught during this field study.

### 2.7. Simpson index of heterogeneity

The fish specimens from artisanal fishermen catches were sorted out according to species, counted and the values fitted into the following model for Simpson index of heterogeneity determination using Simpson index calculator developed by Al Young Studio, 2017 version.

$$D_s = 1 - \sum \frac{\{n_i(n_i-1)\}}{N(N-1)} \text{ where;}$$

$D_s$  = Simpson's index of diversity;

$N$  = Total number of individuals of all species;

$n_i$  = Total number of individuals of the species.

### 2.8. Data analysis

The data generated were analyzed using IBM SPSS version 22.0 (Armonk, NY: IBM Corp) and the results expressed in descriptive statistics by simple percentages and frequencies.

## 3. Results and discussion

The results for the fish diversities in the two water bodies were summarized in Table 1. The families Mormyridae and Bagridae were the dominant in both Goronyo reservoir and the downstream river Rima while one of the rarest occurring families in Goronyo reservoir was Cyprinidae. The species *Mormyrus rume* and *Bagrus bayad macropterus* were the dominant fish species accounting for 23.64%, 11.17% by number and 27.52%, 15.16% by weight in River Rima respectively (Table 2). The same species accounted for 15.31%, 15.60% (by number) and

21.06%, 21.25% (by weight) in Goronyo reservoir. *Lates niloticus* accounted for 0.43% and 10.7% by weight in Rima and Goronyo respectively. *Synodontis membranaceus* occurred rarely (0.11% by number) in River Rima and also accounted for 0.03% and 0.003% by weights in river Rima and Goronyo reservoir respectively.

**Table 1**

The comparative statuses of fish families in numbers, weight and percentage in river Rima and Goronyo reservoir.

Family	Number (A)		Percentage		Weight (kg)		Percentage		Weight	
	RR	GR	RR	GR	RR	GR	RR	GR	RR	GR
Mormyridae	417	291	44.92	26.06	142.72	44.92	54.65	23.22		
Bagridae	157	240	16.92	21.44	65.24	16.92	24.50	39.14		
Claridae	40	47	4.31	4.21	15.32	4.31	6.07	7.76		
Cichlidae	44	98	4.76	8.78	6.93	4.76	2.62	3.44		
Mochokidae	117	123	12.62	11.07	12.44	12.62	4.77	3.23		
Schilbedae	68	93	7	8.32	2.44	5.13	0.93	1.18		
Centropomidae	2	47	0.22	4.21	1.20	58.58	0.47	13.43		
Polypteridae	-	33	-	2.95	-	5.51	-	1.26		
Malapteruridae	11	37	1.19	3.31	1.65	6.64	0.63	1.52		
Lepidoserinidae	22	77	2.26	6.89	4.73	23.40	1.81	5.36		
Clarotidae	10	3	1.08	0.26	2.26	0.67	0.86	0.15		
Characidae	12	7	1.29	0.63	0.23	0.72	1.11	0.17		
Cyprinidae	16	5	1.70	0.45	1.69	0.03	0.65	0.017		
Distichontidae	13	16	1.40	1.43	4.55	5.16	1.72	1.18		
Total	928	1117	100	100	261.06	436.26	100	100		

Keys: RR = River Rima; GR = Goronyo Reservoir; - = Not found; Source: Field work, 2016.

Table 3 showed the results of Simpson index of heterogeneity for the two fairly uniform water bodies (0.92) in both river Rima and Goronyo reservoir. Furthermore, the arrays of fishing gears used by the fishermen are presented in Table 4. Hook and line and gillnets were the most common gears used and accounted for 29.87% and 35.60% respectively. The least (0.31%) gear used by the fishermen at Goronyo reservoir was seine nets. Gourds accounted for 98.28% of fishing equipment at Goronyo reservoir.

**Table 4**

The percentage fishing gears and equipment used by the fishermen at river Rima and Goronyo reservoir.

S/No.	Gear type	Status RR	GR	River Rima (%)	Goronyo (%)
1	Malian trap	X	x	26.74	22.78
2	Gill net	X	x	14.15	35.60
3	Clap net	X	x	23.58	10.85
4	Cast net	X	-	3.14	-
5	Seine net (taru)	X	x	0.31	0.43
6	Hook and line	X	x	29.87	29.28
7	Pole and line	X	x	1.90	0.87
8	Ndrutu	-	x	-	10.85
9	Fish barrier	X	-	0.16	-
10	Yawa	X	-	0.16	-
	Total			100	100
<b>Equipment</b>					
1	Gourd	X	x	96.10	98.28
2	Canoe	-	x	5.90	1.72
	Total			100	100

Source: Field work, 2016.

**Table 3**

The Simpson index of heterogeneity (1 –D) for river Rima and Goronyo reservoir.

Species name	No. of			No. of		
	Individuals (N)	(N-1)	n <sub>i</sub> (n - 1)	Individual (N)	(N- 1)	n <sub>i</sub> (n - 1)
	River	Rima		Goronyo	Reservoir	
<i>Bagrus bayad</i>	216	215	46440	225	224	50400
<i>Bagrus docmac</i>	23	22	506	101	100	10100
<i>Bagrus filamentosus</i>	5	4	20	3	2	6
<i>Auchenoglanis occidentalis</i>	7	6	42	2	1	2
<i>Auchenoglanis biscutatus</i>	5	4	20	2	1	2
<i>Clarotes laticep</i>	24	23	552	4	3	12
<i>Schilbe mystus</i>	53	52	2756	79	78	6162
<i>Schilbe intermedius</i>	68	67	4556	17	16	272
<i>Parailia pelucida</i>	15	14	210	14	13	182
<i>Synodontis clarias</i>	41	40	1640	45	44	1980
<i>Synodontis sorex</i>	14	13	182	9	8	72
<i>Synodontis nigrata</i>	11	10	253	30	29	870
<i>Synodontis membranaceus</i>	2	1	2	3	2	6
<i>Synodontis gobroni</i>	----	----	----	7	8	56
<i>Synodontis budgetti</i>	13	12	156	2	1	2
<i>Synodontis eupterus</i>	15	14	210	---	----	----
<i>Synodontis courteti</i>	10	9	90	6	5	30
<i>Synodontis vermiculatus</i>	15	14	210	---	----	----
<i>Synodontis filamentosus</i>	7	6	42	---	----	----
<i>Clarias anguillaris</i>	12	11	132	17	16	272
<i>Clarias gariepinus</i>	24	23	552	30	29	870
<i>Hetrobronchus bidorsalis</i>	7	6	42	--	----	----
<i>Alestes baremose</i>	12	11	132	5	4	20
<i>Alestes nurse</i>	4	3	12	---	----	----
<i>Hydrocinus forskalii</i>	2	1	2	---	----	----
<i>Malapterurus electricus</i>	11	10	110	37	36	1332
<i>Lates niloticus</i>	47	46	2162	2	1	2
<i>Oreochromis niloticus</i>	42	41	1722	94	93	8742
<i>Hemichromis fasciatus</i>	2	1	---	---	----	----
<i>Tilapia zilli</i>	----	----	----	2	1	2
<i>Saratherodon melanotheron</i>	2	1	2	2	1	2
<i>Mormyrus rume</i>	279	278	77562	206	205	42230
<i>Marcusinius senegalensis</i>	57	56	3192	16	15	240
<i>Gnathemona abadii</i>	28	27	756	38	37	1406
<i>Hyperopisus bebe occidentalis</i>	130	129	16770	98	97	9506
<i>M. tapirus</i>	10	9	90	27	26	702
<i>Petrocephalus bovie</i>	98	97	9506	16	15	240
<i>Mormyrus macrophthalmus</i>	--	---	----	18	17	306
<i>Labeo senegalensis</i>	12	11	132	9	8	72
<i>Labeo coubie</i>	3	2	6	---	----	----
<i>Barbus occidentalis</i>	--	---	----	4	3	12
<i>Protopterus annectens</i>	21	20	420	77	76	5852
<i>Polypterus senegalus senegalus</i>	---	---	----	3	2	6
<i>Distichodus rostratus</i>	13	12	156	16	15	240

<i>Distichodus engycephalus</i>	2	1	2	---	--	---
Σ of the Parameters			1322			1232
Total of Fish Species	40			35		
Total number of fish Specimens	1362			1224		
Average Population Size	34.06			34.07		
Region	Rima			Goronyo		
Simpson Index of heterogeneity (1 -D)			0.90			0.90

Source: Field work, 2016.

**Table 2**

The fish species composition, relative abundance and dominance in both Goronyo Reservoir and River Rima.

Family/Species	Status		Number (A)		Weight (kg)		Percentage (RA)		Percentage (%)		Weight		Remarks	
	RR	GR	RR	GR	RR	GR	RR	GR	RR	GR	RR	GR	RR	GR
<b>Mormyridae</b>														
<i>Mormyrus rume</i>	x	X	218	172	77.05	60.19	23.64	15.31	27.52	21.06	D	D		
<i>M. tapirus</i>	x	X	16	17	6.04	8.43	1.74	1.53	2.16	1.55	S	S		
<i>M. macrophthalmus</i>	x	X	3	6	0.83	1.8	0.33	0.54	0.30	0.33	O	O		
<i>Hyperopisus bebe occidentalis</i>	x	X	90	47	33.31	11.74	9.76	4.24	11.90	2.16	D	S		
<i>Marcusinius senegalensis</i>	x	X	52	14	5.23	0.67	5.64	1.26	1.87	0.12	S	O		
<i>G. abadii</i>	x	X	38	25	24.95	18.30	4.12	2.25	8.91	3.36	S	S		
<i>Petrocephalus bovie</i>	x	X	98	10	-	0.20	----	0.90	---	0.04	E	R		
Sub total			417	291	147.41	101.33	45.19	25.23	52.65	18.62				
<b>Bagridae</b>														
<i>Bagrus bayad macropterus</i>	x	X	103	173	42.43	115.60	11.17	15.60	15.16	21.25	D	D		
<i>Bagrus docmac</i>	x	X	25	60	16.82	48.49	2.71	5.41	6.01	8.91	S	S		
<i>B. filamentosus</i>	x	X	5	3	1.65	1.60	0.54	0.27	0.58	0.29	O	O		
<i>Auchenoglanis biscutatus</i>	x	X	5	2	0.87	0.5	0.54	0.18	0.31	0.09	O	R		
<i>A. Occidentalis</i>	x	X	7	4	1.32	0.17	0.76	0.36	0.47	9.03	O	S		
Sub total			145	242	63	166.19	15.80	20.82	22.53	40.03				
<b>Clarotidae</b>														
<i>Clarotes laticep</i>	x	X	24	4	4.34	0.70	2.60	0.36	1.55	0.13	S	O		
Sub total			24	4	4.34	0.70	2.60	0.36	1.55	0.13				
<b>Claridae</b>														
<i>Clarias gariepinus</i>	x	X	24	30	7.12	23.44	2.60	2.70	2.54	4.31	S	S		
<i>C. anguillaris</i>	x	X	12	17	5.65	10.44	1.30	1.53	2.02	1.92	S	S		
<i>Hetrobronchus bidorsalis</i>	x	-	4	-	2.55	---	0.43	---	0.91	---	R	E		
Sub total			40	37	12.77	36.43	4.33	4.23	5.47	6.23				
<b>Cichlidae</b>														
<i>Oreochromis niloticus</i>	x	X	42	94	6.45	14.09	4.56	8.00	2.30	2.59	S	S		
<i>Hemichromis fasciatus</i>	x	-	2	-	0.38	-	0.22	----	0.14	----	O	E		
<i>Tilapia zilli</i>	-	X	-	2	-	0.5	----	0.18	----	0.09	E	R		
<i>Saratherodon melanotheron</i>	x	X	2	2	-	0.32	----	0.18	----	0.06	E	R		
Sub total			44	98	6.83	14.91	4.78	8.36	2.44	2.74				
<b>Mochokidae</b>														
<i>Synodontis clarias</i>	x	X	41	55	4.56	8.29	4.44	4.96	1.63	1.52	S	S		
<i>S. nigrata</i>	x	X	11	30	1.29	3.12	1.19	2.70	0.46	0.57	O	O		
<i>S. sorex</i>	x	X	14	9	0.88	1.09	1.51	0.81	0.31	0.20	O	O		
<i>S. membranaceus</i>	x	X	2	3	0.09	0.02	0.11	2.27	0.03	0.003	R	R		
<i>S. budgetti</i>	x	X	13	2	0.59	0.69	1.4	0.18	0.21	0.13	O	O		
<i>S. eupterus</i>	x	-	-	15		4.44	--	1.35	----	0.82	E	O		

<i>S. courteti</i>	x	X	10	6	0.27	0.46	1.08	0.54	0.10	0.08	O	O
<i>S. vermiculatus</i>	x	-	5	-	0.08	-	0.54	---	0.01	----	R	E
<i>S. filamentosus</i>	x	-	7	-	0.24	-	0.76	----	0.27	----	O	R
<i>S. gobroni</i>	-	X	7	-	19	-	0.76	----	6.79	----	O	E
Sub total			109	120	31	121.58	11.79	12.60	9.81	3.32		
<b>Schilbedae</b>												
<i>Schilbe mytus</i>	x	X	53	79	1.47	2.82	5.75	7.12	0.53	0.53	S	O
<i>Parailia pelucida</i>	x	X	15	14	0.14	0.11	1.63	1.26	0.05	0.02	S	O
Sub total			68	93	1.61	2.93	7.38	8.38	0.58	0.55		
<b>Centropomidae</b>												
<i>Lates niloticus</i>	x	X	47	2	1.2	58.58	0.22	4.24	0.43	10.77	O	O
Sub total			2	47	1.2	58.58	0.22	4.24	0.43	10.77		
<b>Polypteridae</b>												
<i>Polypterus senegalus senegalus</i>	-	X	-	33	-	5.51	---	2.98	---	1.01	E	S
Sub total			-	33	-	5.51	----	2.98	----	1.01		
<b>Malapteruridae</b>												
<i>Malapterurus electricus</i>	x	X	11	37	1.65	6.64	1.19	3.34	0.59	1.22	S	S
Sub total			11	37	1.65	6.64	1.19	3.34	0.59	1.22		
<b>Lepidoserinidae</b>												
<i>Protopterus annectens</i>	x	X	21	77	4.73	23.40	2.28	6.94	1.69	4.30	S	S
Sub total			21	77	4.73	23.40	2.28	6.94	1.69	4.30		
<b>Characidae</b>												
<i>Alestes baremose</i>	x	X	12	5	1.14	0.03	1.30	0.45	0.41	0.005	S	R
<i>A. Nurse</i>	x	-	4	-	0.55	-	0.43	-	0.20	----	O	E
Sub total			16	5	0.55	0.03	1.73	0.45	0.61	0.005		
<b>Cyprinidae</b>												
<i>Labeo senegalensis</i>	x	X	9	3	0.22	0.55	0.98	0.27	0.08	0.10	O	O
<i>L. coubie</i>	x	-	3	-	0.08	-	0.32	---	0.03	----	R	E
<i>Barbus occidentalis</i>	-	X	-	4	-	0.17	----	0.36	---	0.03	E	R
Sub total			12	9	0.30	0.72	1.3	0.63	0.11	0.13		
<b>Distichontidae</b>												
<i>Distichodus rostratus</i>	x	X	13	16	4.55	5.16	1.4	1.44	1.63	0.95	S	O
Sub total			13	16	4.55	5.16	1.4	1.44	1.63	0.95		
Grand Total			922	1,109	279.94	544.11	100	100	100	100		

Source: Fieldwork, 2016. X = Found, - = not found, RR = River Rima, GR = Goronyo Reservoir, A = Abundance, R.A. = Relative abundance, D = Dominant, S = Subdominant, O = Occasional, R = Rare, E = Extinct.

The results on fish biodiversity in the two water bodies followed a similar pattern reported in some Nigerian water bodies. Rivers are known to typically support more fish species than their associated reservoirs, often as a result of large scale changes in regimes of temperature, turbidity, flow, allochthonous nutrient inputs and availability of food resources (Williams et al., 1998). The result of the present finding had indicated the species compositions were almost equal with 14 versus 15 for River Rima and Goronyo reservoir respectively.

However, fish distribution and abundance in tropical water bodies have been variously attributed to several factors. Some of these factors were principally the water depth (Chapman and Kramer, 1991), water temperature and transparency (Agremier and Karr, 1983), reported availability of food and migratory nature of some fishes (Adebisi, 1988; Wine Miller and Japsen, 1988). Besides, to some of these factors responsible for different fish distribution in some water bodies, the prevailing ecological conditions like the nutrients level also plays role in uneven fish distribution (Ita et al., 1982). In this study, the nutrients level, current speed and nature of the bottom deposit, may have acted singly or synergistically to influence the presence status of fish abundance and distribution in River Rima and Goronyo reservoir. Similar observations were reported by Petr (1975) and Odum (1995) in the Ethiopie River of southern Nigeria.

Ita et al. (1982) reported only nine fish families in the river Rima, but this study observed 14 families, thereby indicating encroachment or inversion and colonizing some portion of the two water bodies by new species. Both the two water bodies exhibited high species richness with 40 and 35 fish species, for river Rima and Goronyo reservoir respectively compared to 20 species (Ita et al., 1982). This is because, of the difference in scope and



length of periods in the two studies. Other studies that reported high species richness in Nigerian water bodies like this study includes the study conducted by Bolarinwa et al. (2015) in the coastal areas of Ondo State, Nigeria. The authors reported 36 families represented by 67 species therefore exhibiting higher species richness. Ali and Abubakar (2015) also reported the presence of 26 species from 14 families in Dadinkowa dam in Gombe, with *Bagrus bayad* been the dominant fish species. A similar study in some parts of Africa was conducted by Mohammed (2012) and reported 23 species from 13 families in Jabel Awlia dam and White Nile River in Sudan with *Alestes* species accounting for 19.26% of the total catches.

Compared to this study, however, some Nigerian water bodies exhibited low species richness; for example, Edward (2012) reported low species richness (8 species from 5 families) in Egbe Reservoir in Ekiti state with Cichlids dominating the catches compared to this study. The dominance of *Mormyrus rume* in terms of number in both River Rima (23.64%) and Goronyo reservoir (15.31%) shows that the dominant fish species as a detritivore is the primary consumers, thus indicating the high potential to fish production under adequate management (Ita and Balogun, 1982).

Most of the gears used by the fishermen were local fishing gears commonly used by Nigerian fishermen, with gillnet and hook and line dominating the rest. Reed et al. (1967) also reported high use of gillnet and hook and lines by the northern Nigerian fishermen. Furthermore, Dan-kishiya et al. (2012) also reported the dominance of gillnets by the fishermen of Lower Usuma reservoir, Bwari in Abuja, Nigeria. This shows the versatility of gill nets in being able to trap fish of different sizes and shapes. According to Sikoki et al. (1998), gill nets could catch fish of all sizes, shapes and species in all water habitats.

#### 4. Conclusion and recommendations

The findings in this study revealed high species richness and diversity in both River Rima and Goronyo reservoir. These findings have provided baseline information for proper fisheries management in the reservoir and the river Rima in terms of species enhancement, management and protection. Based on the results on fish biodiversity in the reservoir and river Rima, the present work recommended the restocking of the water bodies with *Lates niloticus* and *Labeo coubie*, regulating the fishing gears' use and conducting a further study on forage-predator fish species ratio.

#### Acknowledgments

The authors would like to acknowledge the efforts of fishermen of River Rima and Goronyo reservoir who provided the fish specimens for this study. We also thank the management of the Goronyo reservoir for making their facilities accessible for us during the field work.

#### References

- Adebisi, A.A., 1988. Changes in the structural and functional components of fish community of a seasonal river. Arch. Hydrobiol., 113, 457-463.
- Agremier, P.L., Kerr, J.R., 1983. Fish communities along environmental gradients in system tropical streams. Environ. Biol. Fish., 9, 457-463.
- Ali, J., Abubakar, U.M., 2015. Fish species biodiversity and abundance of Dadinkowa Dam, Gombe State, Nigeria. Int. J. Innovat. Res. Dev., 3(6), 374-378.
- Bagenal, T., Tesch, F., 1978. Age and growth, In: Bagenal, T. (Ed.), Methods for assessment of fish production in fresh waters, 3rd Edition, IBP Handbook No. 3, Blackwell Science Publications, Oxford.
- Balogun, J.K., 2005. Fish distribution in a small reservoir: A case study of Kangimi reservoir. J. Appl. Sci. Environ. Manag., 9(1), 93-97.
- Benech, V., Durand, J.R., Quensiere, J., 1983. Fish communities of lake Chad and associated rivers and floodplains. In Carmouze, J.P., Durand, J.R., Lévêque, C. (Eds), Lake Chad: Ecology and productivity of a shallow tropical ecosystem. The Hague: Dr W. Junk. 293-357.
- Bolarinwa, J.B., Fasakin, E.A., Fagbenro, A.O., 2015. Species composition and diversity of the coastal waters of Ondo State, Nigeria. Int. J. Res. Agr. Forest., 2(3), 51-58.

- Chapman, L.J., Kramer, D.L., 1991. Limnological observations of an intermittent tropical dry forest streams. *Hydrobiol.*, 226, 153-166.
- Dan-Kishiya, A.A., Olatunde, A.A., Balogun, J.K., 2012. The status and prospects of artisanal fisheries of lower Usuma reservoir, Bwari, F.C.T. Abuja, Nigeria. *Res.*, 4(2), 4-7.
- Dankishiya, A.S., Olatunde, A.A., Balogun, J.K., 2013. Ichthyofauna composition and diversity of a tropical water reservoir: A case study of lower Usuma reservoir in Bwari Abuja, Nigeria. *Am. J. Res. Comm.*, 1(9), 188-203.
- Edward, J.B., 2012. Evaluation of the fisheries potentials of Egbe reservoir, Ekiti State, Nigeria. *Greener J. Biol. Sci.*, 3(7), 260-267.
- Eniolorunda, N.B., Dankani, I.M., Yusuf, N., 2012. A Remote Sensing (RS) and Geographic Information System (GIS) approach to estimating electric power consumption: A case of Sokoto Metropolis, Sokoto State, Nigeria. *Int. J. Manag. IT Eng.*, 2(9), 231-242.
- Google Earth, 2016. Image viewer, retrieved on 23/01/2016 <https://earth.google.com/web/@9.0338725,8.677457>
- Gupta, S.R., Gupta, P.C., 2008. *General and Applied Ichthyology (Fish and Fisheries)*. S. Chand Publication, New Delhi.
- Gusau, I.A., Bilkisu, A.S., Ipinjolu, J.K., 2016. Artisanal Fisheries of Bakolori Reservoir in Zamfara State, Northwestern Nigeria. *Equity J. Sci. Technol.*, 3(2), 43-50.
- IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp.
- Ibrahim, B.U., Auta, J., Balogun, J.K., 2009. A survey of the artisanal fisheries of Kontagora reservoir, Niger State, Nigeria. *Bayero J. Pure Appl. Sci.*, 2(1), 47-51.
- Ita, E.O., 1993. *Inland Fisheries Resources of Nigeria*, FAO of the United Nation, Rome: CIPA Occasional Paper no. 20.
- Ita, E.O., Balogun, J.K., 1982. Report of pre-impoundment fisheries survey of Goronyo reservoir, Sokoto State, Nigeria. A Report Submitted to Sokoto Rima Basin Development Authority, 86p.
- Ita, E.O., Balogun, J.K., Ademula, A., 1982. A preliminary report of pre-impoundment fisheries survey of Goronyo reservoir, Sokoto State, Nigeria. Technical, Sokoto-Rima Rivers Development Authority, Nigeria, Sokoto.
- Joseph, B.B., Emanuel, A.F., Adedapo, O.F., 2015. Species composition and diversity of the coastal waters of Ondo State, Nigeria. *Int. J. Agr. Forest.*, 2(3), 51-58.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Desertification Synthesis*. World Resources Institute, Washington, DC., USA.
- Mohammed, M.O., 2012. Fishes List of Jabel Awlia Dam Reservoir in the White Nile River, Sudan, *Bulletin of Environment. Pharmacol. Life Sci.*, 1(5), 26-29.
- Odum, O., 1995. Fish Distribution in Ethiopie River, Southern Nigeria. *Trop. Freshwat. Biol.*, 4, 53-64.
- Petr, T., 1975. Some factors associated with the initial high fishes catches in new African man-made lakes. *Arch. Hydrobiol.*, 32-49.
- Reed, W., John, B., Hopson, A.J., Jennings, J., Yaro, I., 1967. *The fish and fisheries of northern Nigeria (1<sup>st</sup> Ed.)* Zaria, Gaskiya Corporation Limited, Zaria, Nigeria.
- Sembenelli Consulting, 2010. *Goronyo Main and Secondary Dam (pdf)* Archived from the original retrieved 2010-05-21.
- Shinkafi, A.S., Ukaola, L.N., Thekir, N.A., 2013. Some aspects of growth and reproduction in Nile Perch (*Lates niloticus*, Linne 1760) from River Rima, Northwestern Nigeria. *Niger. J. Fish. Agr.*, 1(1), 31-41.
- Sikoki, F.D., Hart, A.I., Abowei, J.F.N., 1998. Gill net selectivity and fish abundance in the lower Nun River, Bayelsa State, Nigeria. *J. Appl. Sci. Environ. Manag.*, 1, 13-19.
- William, R.M., Halwart, T., Barig, H., 1998. Integrating fisheries and agriculture to enhance fish production and food security. *FAO Aquaculture Newsletter*, 20, 3-8.
- Wine Miller, K.O., Japsen, D.B., 1998. Effects of seasonality and fish movement on tropical river food webs. *J. Fish. Biol.*, 53(Supplementary A), 267-296.

**How to cite this article:** Gusau, I.A., Muhammad, U., Bilbis, B.D., 2020. A comparative study in fish biodiversity in Goronyo reservoir and downstream river Rima in Sokoto State Northwestern Nigeria. *Agricultural Advances*, 9(7), 545-554.

**Submit your next manuscript to Sjournals Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in DOAJ, and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.sjournals.com](http://www.sjournals.com)

