

Contents lists available at Sjournals



Journal homepage: [www.Sjournals.com](http://www.Sjournals.com)



**Original article**

## **Influence of biologically treated wheat straw diet on *in vitro* rumen fermentation, methanogenesis and digestibility**

**M.S. Mahesh<sup>a,\*</sup>, M. Mohini<sup>a</sup>, D. Kumar<sup>b</sup>, R. Sheel<sup>a</sup>, S.P. Sawant<sup>a</sup>, P. Jha<sup>a</sup>**

<sup>a</sup>*Dairy Cattle Nutrition Division, National Dairy Research Institute (Deemed University), Karnal, Haryana-132 001, India.*

<sup>b</sup>*Animal Nutrition Division, Indian Veterinary Research Institute, Izatnagar, U.P., India.*

\*Corresponding author; Dairy Cattle Nutrition Division, India; Tel: +91-9541853329.

### ARTICLE INFO

### ABSTRACT

*Article history:*

Received 20 June 2013

Accepted 25 June 2013

Available online 29 June 2013

*Keywords:*

Digestibility

Gas production

Microbial protein synthesis

Methane

Wheat straw

The present study was conducted to assess the nutritive worth of diets comprising of wheat straw (T1 diet) and basidiomycetes white-rot fungal (WRF) isolate RCK-SC treated wheat straw (T2 diet) for *in vitro* gas production, digestibility and rumen fermentation parameters. Diets (on dry basis) containing 70% straw, 10% berseem green forage, 17% groundnut cake, 2% mineral mixture and 1% salt were formulated. It was observed that *in vitro* total gas (GV 24 h), true DM and OM digestibility (48 h), *in vitro* NDF digestibility (48 h), microbial biomass production (MBP) and short chain fatty acid (SCFA) production were significantly ( $P<0.05$ ) higher in T2 diet to the extent of 9.71, 17.39, 14.92, 17.45, 6.34, 9.34%, respectively when compared to T1. While, total N and NH<sub>3</sub>-N in rumen liquor of both the diet remained same. Volatile fatty acids like propionate concentration increased ( $P<0.05$ ) with a decrease in acetate to propionate ratio (A:P). Methane production expressed as L/kg IVTDMD was significantly ( $P<0.05$ ) lower (13.86%) in T2 diet. This study demonstrates that fungal treatment of wheat straw using basidiomycetes WRF isolate RCK-SC produces better quality straw and its incorporation benefits rumen fermentation *in vitro*.

© 2013 Sjournals. All rights reserved.

## 1. Introduction

One of the reasons for suboptimal performance of livestock in the tropical countries can be the feeding of poor quality dry roughages like straws (Logeswari Monica et al., 2013), which constitute a major part of ruminant ration, including in India. Wheat straw, a major energy-rich cereal straw available in Northern India (up to 109.9 MMT, Saritha et al., 2012) is low in protein and minerals besides the presence of lignin-cell wall polysaccharide complex, which make it inaccessible by rumen microbiota leading to low digestibility (García-Cubero et al., 2012) that in turn limits its fullest nutritional utility in ruminants. Biological treatment by solid state fermentation (SSF) using basidiomycetes white-rot fungi (WRF) has been employed as a means of bioremediation of straws to improve nutritive value by selectively reducing lignin content (Villas-Bôas, 2002; Hassim et al., 2012), which is a non-hazardous and eco-friendly method. Different WRF like *Ceriporiopsis* sp., *Cyathus stercoreus*, *Pleurotus* sp., *Lentinus edodes*, *Phlebia* sp., *Phanerochaete chrysosporium*, *Ganoderma* sp. and *Trametes versicolor* have been used successfully to improve the nutritive value of various substrates viz. wheat straw, paddy straw, madake bamboo, bermuda grass, corn-stalks and oil palm fronds etc. under SSF (Okano et al., 2009; Akinfemi, 2010; Arora and Sharma, 2012; Hassim et al., 2012; Omer et al., 2012; Shrivastava et al., 2012).

Livestock are responsible for 18% of the global anthropogenic greenhouse gas emissions, accounting for about 37% of the total anthropogenic methane emissions (Patra and Yu, 2012). Methanogenesis represents a net loss of 2-12% of ingested gross energy, depending on the quality of feeds fermented in the rumen (Patra, 2012). Ruminants fed on crop residues are reported to emit a large quantity of methane (Zhu et al., 2008) due to longer ruminal residency period (Moss et al., 1994). A few in vitro studies also indicated that fungal treated substrates yield low methane in the rumen (Jalc et al., 1994; Akinfemi, 2010).

As the gas production technique (Menke and Steingass, 1988) is one of the best techniques currently available to study the rate of in vitro organic matter fermentation in rumen fluid (Krishnamoorthy et al., 2005), this technique is employed in the current study.

Objective of the present investigation was to evaluate the nutritional quality in terms of rumen fermentation and digestibility of diet comprising mainly of wheat straw treated with ligninolytic WRF RCK-SC by in vitro methods.

## 2. Materials and methods

### 2.1. SSF of wheat straw

Wheat straw was dried at 60 °C and sieved to attain a uniform particle size (1.5-2.0 cm) before SSF (Shrivastava et al., 2012). The unfermented and fermented straws were ground in the laboratory mill (Remi Motors, Delhi, India) and sieved (30 mesh) for analytical purposes.

The fungus used in the study was isolated from the leaf litter in Delhi ridge forest area and temporarily named as basidiomycetes WRF RCK-SC, maintained on malt extract agar (MEA) in the University of Delhi South Campus, New Delhi. The cultures were stored at 4°C and subcultured every fortnight. Inoculation of wheat straw with fungal inoculums was carried out as per Shrivastava et al. (2012) with a final substrate to moisture ratio of 1:3 and incubated at 30 °C and 60% relative humidity for 10 days. The uninoculated trays without fungal inocula served as control. The inoculated wheat straw trays in triplicates were harvested at regular intervals and weight loss of wheat straw was determined by deducting the weight of oven dried fermented straw (at 60 °C to a constant weight) from the weight of dried control straw (Shrivastava et al., 2011).

### 2.2. Experimental diets

The two experimental diets were prepared as below considering maximum inclusion of wheat straw.

T1 diet: 70% wheat straw (WS) + 10% berseem fodder + 17% groundnut cake + 2% mineral mixture + 1% salt.

T2 diet: 70% fungal treated wheat straw (FT-WS) + 10% berseem fodder + 17% groundnut cake + 2% mineral mixture + 1% salt.

### 2.3. Proximate, cell wall analysis, in vitro gas production, methane and digestibility

Proximate and cell wall analysis were performed according to AOAC (2005) and Van Soest et al. (1991), respectively. Hemicellulose was the difference between neutral detergent fibre (NDF) and acid detergent fibre

(ADF). Acid detergent lignin (ADL) was assayed by solubilizing ADF with 72% (w/w) sulfuric acid (Van Soest et al., 1991). Cellulose was calculated by subtracting ADL from ADF. Total carbohydrates (T-CHO) were calculated by subtracting crude protein, ether extracts and ash contents from 100 (Sniffen et al., 1992). All the chemical analyses were in triplicates.

For in vitro gas production study (Menke and Steingass, 1988), rumen fluid was collected from two cannulated Karan-Fries bulls (5-6 years, body weight  $450 \pm 7.3$  kg) fed on maintenance ration. Around  $200 \pm 10$  mg of air equilibrated substrates of both T1 and T2 diets were taken with 30 ml of buffered rumen inoculum (10 ml rumen fluid, 5ml bicarbonate buffer, 5 ml macro-mineral solution, 0.0025 ml micro-mineral solution and 10 ml distilled water) under continuous flushing with CO<sub>2</sub> into 100 ml calibrated syringes (Fortuna optima, Germany) along with blanks. After 24 h of incubation, net gas produced (GP24 h) was recorded from visual assessment of the calibrated scale on the syringe and was corrected for blanks. In vitro microbial biomass production (MBP) was calculated (Blümmel et al., 1997) considering 2.20 as the stoichiometric factor, and in vitro short chain fatty acids (SCFA) were calculated by the equation developed by Getachew et al. (2002), which are as follows.

$$\text{MBP (mg)} = \text{TDOM (mg)} - (\text{GP24 h} \times 2.20)$$

where TDOM is truly digestible organic matter

$$\text{In vitro SCFA (mM/0.2g DM)} = (0.0239 \times \text{GP24 h}) - 0.0601$$

For in vitro true dry matter digestibility (IVTDMD), separate incubations (48 h) of the 2 diets were carried out (Menke and Steingass, 1988) and, IVTDMD and in vitro NDF digestibility (IVNDFD) was calculated after recovering syringe contents with neutral detergent solution (Blümmel and Becker, 1997). Upon ashing the residue at 550 °C, in vitro true organic matter digestibility (IVTOMD) was estimated.

For methane estimation, 1 ml gas was sampled from the head space of syringe in an air tight syringe (Hamilton) and injected into gas chromatograph (Nucon 5700, India) equipped with stainless steel column packed with Porapak-N and flame ionization detector (FID). Methane volume (ml) was calculated as methane (%) × total gas produced (ml) in 24 h as suggested by Bhatta et al. (2013) and expressed per unit of IVTDMD and IVTOMD.

**Table 1**

Chemical composition of the experimental diets (% DM basis).

Parameter	T1 diet	T2 diet
Organic matter	91.54	88.51
Crude protein	12.42	13.73
Ether extract	2.36	2.10
Total ash	7.86	10.89
NDF (ash free)	60.33	54.96
ADF (ash free)	41.28	36.94
Hemicellulose	19.05	18.02
Cellulose	32.39	30.32
Acid detergent lignin	9.03	6.76
T-CHO	76.76	72.67

#### 2.4. Rumen fluid analysis

After noting the gas volume at 24 h, contents of each syringe were centrifuged (Heraeus, Germany) at 12,000 × g for 20 min at 4 °C to get clarified rumen liquor, which was used immediately for ammonia nitrogen (NH<sub>3</sub>-N) estimation and preserved at -20 °C for determining individual volatile fatty acids (IVFA) as per Erwin et al. (1961) using gas chromatograph (Nucon 5700, India) equipped with flame ionization detector (FID) and stainless steel column (length 4l; o.d ¼II; i.d 3 mm) packed with Chromosorb-101. Total nitrogen (N) and NH<sub>3</sub>-N were estimated by Kjeldahl method by following only distillation and titration steps.

#### 2.5. Statistical analysis

Data were analyzed using one way analysis of variance (ANOVA) as per Snedecor and Cochran (1994) by Sigmastat software, considering the treatment as a factor with the following model.

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where  $\mu$  = population mean;  $T_i$  = effect of treatment;  $e_{ij}$  = random error.

The final data were tabulated and presented as mean  $\pm$  SE.

### 3. Results

The chemical composition of the experimental diets is presented in Table 1. Results of *in vitro* gas production test (IVGPT) revealed a significant ( $P < 0.05$ ) rise in GP<sub>24 h</sub> in T2 diet. Further, there was an increase ( $P < 0.05$ ) in IVTDMD, IVTOMD and IVNDFD in T2 diet (Table 2). MBP and SCFA were improved significantly ( $P < 0.05$ ) in T2 diet (Table 2). Methane production was lower ( $P < 0.05$ ) in T2 diet compared to T1. Rumen liquor analysis with respect to total N and NH<sub>3</sub>-N showed no difference among the two diets (Table 3). Among the volatile fatty acids, the two diets did not differ significantly ( $P > 0.05$ ) for butyrate, whereas they differed ( $P < 0.05$ ) in acetate and propionate concentration, and acetate to propionate (A:P) ratio.

**Table 2**

Effect of fungal treated wheat straw based diet on *in vitro* GP<sub>24 h</sub>, digestibility (%) and associated parameters.

Parameter	T1 diet	T2 diet
GP <sub>24 h</sub> (ml/g DM)	145.83 <sup>a</sup> $\pm$ 4.41	160.00 <sup>b</sup> $\pm$ 6.29
IVTDMD (%)	47.37 <sup>a</sup> $\pm$ 0.66	55.61 <sup>b</sup> $\pm$ 1.17
IVTOMD (%)	52.0 <sup>a</sup> $\pm$ 0.80	59.76 <sup>b</sup> $\pm$ 0.76
IVNDFD (%)	35.30 <sup>a</sup> $\pm$ 0.39	41.46 <sup>b</sup> $\pm$ 1.02
CH <sub>4</sub> (L/kg IVTDMD)	78.86 <sup>a</sup> $\pm$ 2.19	67.93 <sup>b</sup> $\pm$ 4.89
CH <sub>4</sub> (L/kg IVTOMD)	78.26 <sup>a</sup> $\pm$ 1.99	72.17 <sup>b</sup> $\pm$ 4.41
MBP (mg/g DM)	199.31 <sup>a</sup> $\pm$ 1.74	211.95 <sup>b</sup> $\pm$ 2.47
SCFA (mM/g DM)	3.21 <sup>a</sup> $\pm$ 0.09	3.51 <sup>b</sup> $\pm$ 0.14

<sup>a,b</sup>Means bearing different superscripts within a same row differ significantly ( $P < 0.05$ ).

IVTDMD, IVTOMD, IVNDFD, MBP and SCFA are *in vitro* true dry matter digestibility, *in vitro* true organic matter digestibility, *in vitro* NDF digestibility, microbial biomass production and short chain fatty acid production, respectively.

**Table 3**

Effect of fungal treated wheat straw diet on *in vitro* rumen fermentation parameters.

Parameter	T1 diet	T2 diet
NH <sub>3</sub> - N (mg/dl)	14.82 <sup>a</sup> $\pm$ 0.22	15.61 <sup>a</sup> $\pm$ 0.15
Total N (mg/dl)	56.37 <sup>a</sup> $\pm$ 0.31	58.31 <sup>a</sup> $\pm$ 0.63
Fractions of VFA (%)		
Acetic acid	64.50 <sup>a</sup> $\pm$ 0.60	60.53 <sup>b</sup> $\pm$ 0.52
Propionic acid	30.41 <sup>a</sup> $\pm$ 0.70	33.25 <sup>b</sup> $\pm$ 0.25
Butyric acid	8.25 $\pm$ 0.63	9.18 $\pm$ 0.08
A:P ratio	2.12 <sup>a</sup> $\pm$ 0.05	1.82 <sup>b</sup> $\pm$ 0.02

<sup>a,b</sup>Means bearing different superscripts within a same row differ significantly ( $P < 0.05$ ).

### 4. Discussion

As better quality of straw results in an increase in productivity of ruminants (Walli, 2009), the present experiment was conducted with the objective of testing the nutritional superiority of WRF treated wheat straw diet over the control. The tested fungus was selectively ligninolytic and reduced the lignin content of the treated straw up to 40.33% (Mahesh et al., 2013), thus availing more carbohydrates for optimal ruminal degradation

(Akinfemi, 2010) which lead to the higher digestibility of T2 diet. Similarly, Shrivastava et al. (2012) documented higher in vitro gas volume coupled with increased organic matter digestibility in wheat straw fermented with *Ganoderma sp. rckk02*. Higher gas production and digestibility in T2 diets were in consistent with several previous reports on different biologically treated substrates (Akinfemi, 2010; Shrivastava et al., 2011; Shrivastava et al., 2012; Omer et al., 2012). Recently, Hassim et al. (2012) observed an increased in vitro apparently rumen degradable carbohydrates of oil palm fronds inoculated with *Ceriporiopsis subvermispora* (3 weeks) and *Lentinula edodes* (9 weeks) up to 13 and 10%, respectively, proving the potential of WRF treatment in enhancing digestibility. MBP depends upon a balanced ruminal supply of ammonia, energy and carbon skeletons for amino acid synthesis (Wanapat et al., 2009) and diets with low structural carbohydrate contents are known to increase the efficiency of MBP (Tas and Susenbeth, 2007) as in T2 diet. Higher MBP implies a maximum substrate fixation into microbial cells leading to higher microbial protein flow to the intestine as absorbable amino acids. Furthermore, higher gas volume is positively correlated to in vitro MBP (Krishnamoorthy et al., 1991) which substantiates the present findings. A higher SCFA synthesis reflects an active fermentation of carbohydrates, and a high gas production is expected to yield higher value of SCFA (Blümmel and Ørskov, 1993) as observed in our study for T2 diet. Akinfemi (2010) also recorded a significantly higher SCFA ( $\mu\text{M}$ ) in *Pleurotus ostreatus* (0.928) and *P. pulmonaris* (0.896) treated peanut husks as compared to untreated husk (0.370). The present study shows a positive relationship between GP24 h, IVTOMD and in vitro SCFA synthesis (Menke and Steingass, 1988). Further, two diets differed with respect to in vitro methane production. Reduced methane from fungal treated straw diets could be ascribed to the indirect effect of improved fibre digestion (Sallam et al., 2007) with concurrent availability of fermentable nutrients (Mohini and Singh, 2001).

Ruminal  $\text{NH}_3\text{-N}$  concentration is an index of the extent of protein degradation, and the observed values were above the minimum level required for microbial growth (5–8 mg/dl) in the rumen (Satter and Slyter, 1974), which is in agreement with Tripathi et al. (2008). Pattern of volatile fatty acids in the rumen generally follow the trend of feed digestion in the rumen (Patra and Saxena, 2009), which differed among the two diets. This is because, fungal treated straw contained microbial protein in addition to those found within the straw, whose fermentation in the rumen probably yielded more propionate, and thus decreasing the ratio of A:P making ideal fermentation pattern as propionate is the principal glucose precursor in ruminants (over 75% of total blood glucose; Brockman, 1993). Similar VFA patterns were also noted by Karunanandaa and Varga (1996) in rice leaves treated with different white-rots.

## 5. Conclusion

This study demonstrated that fungal treatment of wheat straw using basidiomycetes WRF RCK-SC for 10 days produces better quality straw compared to untreated one as evidenced through IVGPT and associated parameters. Further, enhanced digestibility observed in treated straw by SSF proves its potential in ruminant nutrition, which needs further investigation through feeding trials involving different categories of ruminants.

## Acknowledgement

Dr. Mahesh sincerely acknowledges Indian Council of Agricultural Research (ICAR) for awarding Junior Research Fellowship (JRF) and Director, NDRI for providing facilities to carry out this study. Thanks are also extended to the staff of the Department of Microbiology, University of Delhi South Campus, New Delhi, for supplying fungal treated wheat straw for the study.

## References

- Alexande, G., 1964. Studies on the placenta of sheep (*Ovis aries* L.). Placental size. *J. Reprod. Fert.* 7, 289-305.
- Akinfemi, A., 2010. Bioconversion of peanut husk with white-rot fungi: *Pleurotus* *treatus* and *Pleurotus* *pulmonarius*. *Livest. Res. Rural Develop.* 22, 3.
- AOAC, 2005. Official Methods of Analysis, 18th ed. Association of Official Analytical Chemists, Washington, DC.
- Arora, D.S., Sharma, R.K., 2009. Enhancement in in vitro digestibility of wheat straw obtained from different geographical regions during solid state fermentation by white-rot fungi. *BioResources* 4, 909–920.

- Bhatta, R., Baruah, L., Saravanan, M., Suresh, K.P., Sampath, K.T., 2013. Effect of medicinal and aromatic plants on rumen fermentation, protozoa population and methanogenesis in vitro. *J. Anim. Physiol. Anim. Nutr.* 97, 446–4556.
- Blümmel M., Becker, K., 1997. The degradability characteristics of 54 roughages and neutral detergent fibre as described by gas production and their relation to voluntary feed intake. *Br. J. Nutr.* 77, 757–768.
- Blümmel, M., Ørskov, E.R., 1993. Comparison of gas production and nylon bag degradability of roughages in predictions feed intake in cattle. *Anim. Feed Sci. Technol.* 40, 109.
- Blümmel, M., Makkar, H.P.S., Becker, K., 1997. In vitro gas production: a technique revisited. *J. Anim. Physiol. Anim. Nutr.* 77:24–34.
- Brockman, R.P., 1993. Glucose and short-chain fatty acid metabolism. In: Forbes, J.M., France, J. (Eds.), *Quantitative Aspects of Ruminant Digestion and Metabolism*. CABI Publ., Cambridge, MA, USA, 249–265 pp.
- Erwin, E.S., Marco, G.J., Emery, E.M., 1961. Volatile fatty acid analysis of blood and rumen fluid by gas chromatography. *J. Dairy Sci.* 44, 1768–1771.
- García-Cubero, M.T., Palacín, L.G., González-Benito, G., Bolado, S., Lucas S., Coca, M., 2012. An analysis of lignin removal in a fixed bed reactor by reaction of cereal straws with ozone. *Bioresour. Technol.* 107, 229–234.
- Getachew, G., Makkar, H.P.S., Becker, K., 2002. Tropical browse: contents of phenolic compounds, in vitro gas production and stoichiometric relationship between short chain fatty acid and in vitro gas production. *J. Agric. Sci. (Camb)* 139, 341.
- Hassim H.A., Lourenc, M., Goh Y.M., Baars J.J.P., Fievez, V., 2012. Rumen degradation of oil palm fronds is improved through pre digestion with white rot fungi but not through supplementation with yeast or enzymes. *Can. J. Anim. Sci.* 92, 79–87.
- Jalc, D., Zitnan, R., Nerud, F., 1994. Effect of fungus treated straw on ruminal fermentation in vitro. *Anim. Feed Sci. Technol.* 46, 131–141.
- Karunanandaa, K., Vargaa, G.A., 1996. Colonization of crop residues by white-rot fungi: cell wall monosaccharides, phenolic acids, ruminal fermentation characteristics and digestibility of cell wall fiber components in vitro. *Anim. Feed Sci. Technol.* 63, 273–288.
- Krishnamoorthy, U., Rymer, C., Robinson, P.H., 2005. The in vitro gas production technique: Limitations and opportunities. *Anim. Feed Sci. Technol.* 123–124, 1–7.
- Krishnamoorthy, U., Steingass, H. and Menke, K.H., 1991. Preliminary observation on the relationship between gas production and microbial protein synthesis in vitro. *Arch. Anim. Nutr.* 5, 521–526.
- Logeswari Monica, C., Tirumurugaan K.G., Vijayarani K., Chandrasekaran C., Kumanan, K., 2013. Bioremediation: An alternate utility of cloned lignin peroxidase from *Phanerochaete chrysosporium*. *Indian J. Anim. Sci.* 83, 506–511.
- Mahesh, M.S., Mohini, M., Sawant, S.P., Jha, P., Kundu, S.S., Kuhad, R.C., 2013. Nutritional evaluation of wheat straw treated with basidiomycetes fungal isolate (RCK-SC) in Sahiwal calves. *Livest. Sci.* (in press).
- Menke, K.H., Steingass, H., 1988. Estimation of the energetic feed value obtained from chemical analysis and in vitro gas production using rumen fluid. *Anim. Res. Dev.* 28:7–55.
- Mohini, M., Singh, G.P., 2001. Methane production on feeding jowar fodder based ration in buffalo calves. *Indian J. Anim. Nutr.* 18, 204–209.
- Moss, A.R., Givens, D.I., Garnsworthy, P.C., 1994. The effect of alkali treatment of cereal straws on digestibility and methane production by sheep. *Anim. Feed Sci. Technol.* 49, 245–259.
- Okano, K., Ohkoshi, N., Nishiyama, A., Usagawa, T. and Kitagawa, M., 2009. Improving the nutritive value of madake bamboo, *Phyllostachys bambusoides*, for ruminants by culturing with the white-rot fungus *Ceriporiopsis subvermispora*. *Anim. Feed Sci. Technol.* 152, 278–285.
- Omer, H.A.A., Ali, F.A.F., Gad, S.M., 2012. Replacement of clover hay by biologically treated corn stalks in growing sheep rations. *J. Agri. Sci.* 4, 257–268.
- Patra, A.K., 2012. Enteric methane mitigation technologies for ruminant livestock: a synthesis of current research and future directions. *Environ. Monit. Assess.* 184, 1929–1952.
- Patra, A.K., Saxena, J., 2009. The effect and mode of action of saponins on the microbial populations and fermentation in the rumen and ruminant production. *Nutr. Res. Rev.* 22, 204–219.
- Patra, A.K., Yu, Z., 2012. Effects of essential oils on methane production and fermentation by, and abundance and diversity of, rumen microbial populations. *Appl. Env. Microbiol.* 78, 4271–4280.

- Sallam, S.M.A., Nasser, M.E.A., El-Waziry, A.M., Bueno, I.C.S., Abdalla, A.L., 2007. Use of an in vitro ruminant gas production technique to evaluate some ruminant feedstuffs. *J. Appl. Sci. Res.* 3, 33–41.
- Saritha, M., Arora, A., Lata, 2012. Biological pretreatment of lignocellulosic substrates for enhanced delignification and enzymatic digestibility. *Indian J. Microbiol.* 52, 122–130.
- Satter, L.D., Slyter, L.L., 1974. Effect of ammonia concentration on microbial production in vitro. *Br. J. Nutr.* 32, 194–201.
- Shrivastava, B., Nandal, P., Sharma, A., Jain, K.K., Khasa, Y.P., Das, T.K., Mani, V., Kewalramni, N.J., Kundu, S.S., Kuhad, R.C., 2012. *Bioresour. Technol.* 107, 347–351.
- Shrivastava, B., Thakur, S., Khasa, Y.P., Gupte, A., Puniya, A.K., Kuhad, R.C., 2011. White-rot fungal conversion of wheat straw to energy rich cattle feed. *Biodegradation* 22, 823–831.
- Snedecor G.W., Cochran, W.G., 1994. *Statistical Methods*. Oxford and IBH Publications, New Delhi.
- Sniffen, C.J., O' Connor, J.D., Van Soest, P.J., Fox, D.G., Russell, J.B., 1992. A net carbohydrate and protein system for evaluating cattle diets. II. Carbohydrate and protein availability. *J. Anim. Sci.* 70, 3562–3577.
- Tas, B. M., Susenbeth, A., 2007. Urinary purine derivatives excretion as an indicator of in vivo microbial N flow in cattle: A review. *Livest. Sci.* 111, 181–192.
- Tripathi, M.K., Mishra, A.S., Misra, A.K., Vaithyanathan, S., Prasad, R., Jakhmola, R.C., 2008. Selection of white-rot basidiomycetes for bioconversion of mustard (*Brassica campestris*) straw under solid-state fermentation into energy substrate for rumen micro-organism. *Lett. Appl. Microbiol.* 46, 364–370.
- Van Soest, P.J., Robertson, J.B., Lewis, B.A., 1991. Methods of dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74, 3583–3597.
- Villas-Bôas, S.G., Esposito, E., Mitchell, D.A., 2002. Microbial conversion of lignocellulosic residues for the production of animal feeds. *Anim. Feed Sci. Technol.* 98, 1–12.
- Walli, T.K., 2009. Nutritional strategies for ruminant production in tropics: Indian context. In: Pattanaik, A.K., Verma, A.K., Kamra, D.N. and Sharma, K. (Eds.). *Animal Nutrition: Preparedness to Combat Challenges*. Proceedings of Animal Nutrition Association World Conference, February 14–17, 2009, New Delhi, India, 49–53 pp.
- Wanapat, M., Polyorach, S., Boonnop, K., Mapato, C., Cherdthong, A., 2009. Effects of treating rice straw with urea or urea and calcium hydroxide upon intake, digestibility, rumen fermentation and milk yield of dairy cows. *Livest. Sci.* 125, 238–243.
- Zhu, W.Y., Liu, J.X., Chang, Y.F., Ibal, M.F., Mao, S.Y., 2008. Rumen methanogenesis and nutritional approaches to the mitigation of ruminant methane Invited paper presented at 13th AAAP, International Symposium, Hanoi, Vietnam. 33–40 pp.