

Contents lists available at [Sjournals](http://Sjournals.com)Journal homepage: www.Sjournals.com**Original article****Impact of maternal effects on ranking of animal models for genetic parameter estimation for birth weight in male Afrikaner cattle in Zimbabwe****N. Assan****Department of Agriculture, Faculty of Science, Zimbabwe Open University Bulawayo Region, Zimbabwe*

*Corresponding author; Department of Agriculture, Faculty of Science, Zimbabwe Open University Bulawayo Region, Box 3550, Bulawayo, Zimbabwe

ARTICLE INFO**ABSTRACT***Article history:*

Received 13 May 2012

Accepted 15 June 2012

Available online 07 July 2012

Keywords:

Variance components

Animal model

Birth

Male

Afrikaner cattle

Variance components for additive direct, additive maternal, permanent environmental maternal effects, the covariance between additive direct and maternal effects were estimated by restricted maximum likelihood, fitting four animal models from 1359 pedigree male birth weight records of Afrikaner cattle of Zimbabwe. All investigated models included a random direct genetic effect, but different combinations of random maternal genetic and permanent environmental effects as well as direct-maternal genetic covariance. The direct heritability (h^2_a) ranged from 0.40 to 0.43 when the maternal genetic effects were included in the model, whereas h^2_a was highest 0.45 when maternal effects were excluded. The maternal heritability (h^2_m) was 0.00 when only maternal genetic effects were included in the model and were 0.09 and 0.09 when the permanent environmental effect of the dam was added. The permanent environmental effect of the dam was negligible. Weak negative covariances between direct and maternal genetic effects (σ^2_{am}) was observed in model which had all the random effects fitted and proved to be the appropriate among the four models using the log likelihood ratio test.

© 2012 Sjournals. All rights reserved.

1. Introduction

The Afrikaner is a predominant cattle breed in southern Africa (Oliver, 1983) and was developed as a draft animal with little attention being given to fertility. Numerous studies have investigated the importance of maternal

effects for pre weaning weight in a mixed model setting for various livestock species (Maniatis and Pollot, 2003). Birth is an early measurable trait which is influenced by the individual's genotype and environment and by the maternal ability of the dam, which is also influenced by the genotype and environment each is a source of variation which has been studied independently (Burferning and Kress). Birth weight could be utilized to choose among populations or a criteria preliminary selection in a given population. For birth weight maternal breeding values takes into account uterine environment which has been found significantly affecting birth weight (Joubert and Hammond, 1958). Ferrell (1991) showed that uterine environment can both restrict and allow for optimum growth of a fetus.

An animal model used to estimate maternal effects typically include direct and maternal genetic effects and maternal permanent environmental effects and considers the covariance between direct and maternal effects (Diop et al 1999; Waldron et al 1993). Genetic models including maternal effects and the covariance of direct and maternal genetic effect, fit data better than simple additive model (Dodenhoff et al 1999). However Meyer (1992b) pointed out that such model may be suboptimal and need to be improved. Although selection schemes for early growth that ignore maternal effects are likely to produce suboptimal genetic response (Gerstmayr, 1992). High negative correlations between direct and maternal genetic effects for early weight trait are common but biologically impossible (Meyer, 1994). Therefore the interpretation of results is not straight forward, reflecting the difficulty in separating these effects from the permanent environmental components when all are estimated in the same analysis (Maniatis and Pollot, 2003). The confusion between these effects leads to high sampling errors and strong sampling correlations between the estimates and often the quantity and quality of data available are insufficient to separate them adequately (Meyer, 1992a).

2. Materials and methods

2.1. Study location

Matopos Research Station ($20^{\circ} 23' S$, $31^{\circ} 30' E$) situated 30 km South West of Bulawayo in Zimbabwe. Altitude is low (800m) and the area experiences low erratic rainfall of less than 450 per annum (Homann et al., 2007). Very high summer temperatures, maximum and minimum mean temperatures of hottest months are $21.6^{\circ} C$ and $11.4^{\circ} C$, respectively with possibility of severe droughts (Hagreveas et al., 2004). The most common type of vegetation is sweet veldt with comparatively high nutritional value of browse and annual grass species (Ward et al 1979). Managed well the rangelands should be able to meet the nutritional requirements of goats (Ncube, 2005) and other livestock (Van Rooyen et al., 2004). However, significant proportion of the rangeland are now degraded, resulting in low biomass and thus limited feed resource of poor quality particularly during the dry season (Hlatshwayo, 2007). Day et al (2003) and Gambiza and Nyama (2000) give a detailed description of the climate and vegetation type, respectively. Herd history and management of the herd were described by Assan (2006).

2.2. Herd history and management

In 1957 two lines of grade Afrikaner cows each were established from a common gene pool and subjected to different nutritional and regimes, termed environments. The supplemented line cows were offered supplements during the dry season (9 kg silage and 1 kg cotton seed meal per head per day) and were mated to calve early (October to December) relative to the expected onset of rains. The non supplemented line cows were mated to bulls two months later than those in supplemented line. Selection within line was based on weight at weaning for bulls and weight prior to mating for replacement heifers. Cows were culled for poor productivity (infertility and low calf weaning weight)

Routine cattle veterinary practices were followed. Cows were naturally bred and breeding season was limited to 90 days period from 1 January each year. Single sire herds comprised of one bull to 30 females were introduced to the breeding herd for mating when they had attained two years of age and bulls were seldom used for service until they were three years old. Calves were numbered by means of ear tag and were weighed and recorded within 18 hours of birth. At the same time both the calf number and that of the sire were recorded thereafter all calves weighed and weaned at an average age of 210 days.

2.3. Statistical methods

Data on birth weight in male Afrikaner cattle were obtained from Matopos Research Station, Bulawayo, Zimbabwe. The data included a total of 1359 male's pedigree birth weight records from 32 sires and 346 dams of male Afrikaner cattle of Zimbabwe. Genetic parameters were estimated using the Average Information Restricted Maximum Likelihood (AIREML) methodology (Gilmour, 1995) using an Animal Model. The analytical models included fixed effects of year of birth and age of dam. Fixed factors for models for all traits were determined through preliminary analyses using procedure GLM of SAS (1996) (SAS Inst. Inc., Cary, NC). Fixed factors (main effects and interactions) and covariates were tested and removed from the model if found non-significant ($P > 0.01$), with non-significant effects rejected sequentially.

A simplex algorithm is used to search for variance components to minimize the function, $-2\log$ likelihood (L). Convergence was assumed when the variance of the function values ($-2\log L$) of the simplex was $<10^{-8}$. A log likelihood ratio test was used to choose the most suitable random effects model for post weaning growth. The reduction in $-2 \log L$ when a random effect was added to the model was calculated. If this reduction was greater than the value of the Chi-square distribution with one degree of freedom ($p < 0.05$) the additional random effect fitted was considered significant. When log likelihood did not differ significantly ($p > 0.05$), the model that had the fewer number of parameters was selected as the most appropriate. Four different animal models were fitted for male birth weight by ignoring or including maternal genetic effect, covariance between direct-maternal effects, maternal environmental effect that the four different models were:

Model 1 was a model with animal additive genetic effects and maternal genetic effect as the random effect other than the residuals:

$$y = Xb + Z_a a + Z_m m + e \quad (1)$$

Model 2 included random effect of permanent maternal environment

$$y = Xb + Z_a a + Z_c c + e \quad (2)$$

Model 3 included both permanent environmental and genetic maternal effects but did not allowed correlation between direct and maternal genetic random effects.

$$y = Xb + Z_a a + Z_m m + Z_c c + e \quad \text{Cov}(a,m) \neq A\sigma_{am} \quad (3)$$

Model 4 was the same as model 3 but assumed correlation between direct and maternal genetic random effects.

$$y = Xb + Z_a a + Z_m m + Z_c c + e \quad \text{Cov}(a,m) = A\sigma_{am} \quad (4)$$

where Y is the vector of observations b , a , m , c and e are the vectors of fixed effects, direct additive genetic effects (animal), maternal genetic effects, permanent environmental effect of dam and the residual, respectively. X , Z_a , Z_m , and Z_c , are the incidence matrices of fixed effects, direct additive genetic effects, maternal genetic effects and permanent environmental effect of dam. (Co) variances were described as: $V(a) = \sigma_a^2 A$, $V(m) = \sigma_m^2 A$, $V(c) = \sigma_c^2 I$, $V(e) = \sigma_e^2 I$ and $\text{Cov}(a,m) = \sigma_{am} A$, where σ_{am} is the covariance between direct and maternal genetic effects, σ_a^2 the direct additive genetic variance, σ_m^2 the maternal genetic variance, σ_c^2 the variance of the permanent environmental effect of the dam and σ_e^2 the variance of the residuals. A is the numerator relationship matrix between animals, I the identity matrix. In the present study the fixed part of the model included age at slaughter, sex and year of slaughter of progeny. Heritability of total additive genetic contribution to a maternally influenced trait was calculated according to the following equation (Willham, 1972).

$$h_T^2 = \frac{\sigma_a^2 + 0.5\sigma_m^2 + 1.5\sigma_{am}^2}{\sigma_p^2}$$

3. Results and Discussion

Means, standard deviation and characteristics of the data structure and covariance components are summarized in Table 1 and 2. It was noted that the relative values of direct heritability were influenced by the model used in the analysis (Table 2), which may emphasize the need of ranking of models for appropriateness in genetic evaluation of birth weight in male Afrikaner cattle. Comparison of Model 1 and 2 showed that exclusion of permanent environment maternal effects slightly increased the direct heritability by 5%, however the direct heritability decreased by 7% when both maternal genetic and permanent environmental effects were take into account.

Table 1

Structure and descriptive statistics of the data set of male Afrikaner cattle of Zimbabwe.

Item	Magnitude
Pedigree records	1359
No base parents	161
No animals	2266
No sires	32
No dams	346
Mean (kg)	30.54
Coefficient of Variation (%)	13.30
Standard Deviation (kg)	4.86
Coefficient of Determination (%)	0.50

Table 2

Estimates of covariance components and genetic parameters of birth weight in male Afrikaner cattle of Zimbabwe.

Item	Model 1	Model 2	Model 3	Model 4
σ_a^2	22.11	23.11	21.14	21.40
σ_m^2	0.45		4.77	4.80
σ_{am}^2				-0.27
σ_{pe}^2		0.80	0.36	0.14
σ_e^2	28.85	28.01	26.46	26.25
σ_p^2	51.41	51.93	52.74	52.32
h_a^2	0.43	0.45	0.40	0.41
h_m^2	0.00		0.09	0.09
r_{am}				-0.003
c^2		0.02	0.01	0.00
h_T^2	0.43	0.45	0.45	0.36
-2Log L	15574	15563	15350	15248

σ_a^2 = direct additive genetic

σ_m^2 = maternal additive genetic variance

σ_{am}^2 = direct and additive variance

σ_{pe}^2 = permanent environmental dam variance

σ_p^2 = phenotypic variance = sum of variance and covariance components

σ_e^2 = error variance

h_a^2 = direct heritability

h_m^2 = maternal heritability

r_{am} = direct and maternal genetic correlation

h_T^2 = total heritability (total genetic effect)

Estimates of relative variance due to maternal permanent environmental effects decreased by 55% when both maternal genetic and permanent environmental effects were included in Model 3, and decreased by 83% when a genetic correlation between direct and maternal effects was assumed. The log likelihood showed a better fit with the full model (Model 4). In agreement with the review of Robison (1981), on the whole estimates of heritability and relative values due to permanent environmental effects for Model 3 were well within the range of comparable literature values. Koch et al., (1973) reported that genetic and permanent environmental component of maternal ability and covariance of individual and maternal effects account for 15 to 20 % of variation in birth weight which may be closer to our observation of 10%. The present study indicates that the maternal component for birth weight in the population is much of maternal genetic than maternal environmental. Inclusion of a covariance between direct and maternal genetic effects in Model 4 did not change the magnitude of maternal genetic effects. The range of maternal heritability of birth weight in male Afrikaner cattle in the present study were

within the range reported by several authors. Comparable maternal heritability estimates range of 0.05 – 0.14 were reported in literature (Bertrand et al., 1987; Trus et al., 1988; Meyer, 1992). Brown et al., (1969); Nelsen et al., (1984); Cantet et al., (1988) and Trus et al., (1988) reported higher birth weight maternal heritability estimates of 0.30, 0.82, 0.27 and 0.42, respectively in beef cattle. The maternal heritability estimates were lower than direct heritability which coincides with the findings of several authors (Meyer, 1992; Brown et al., 1996; Trus et al., 1988) however in contrary with what was reported elsewhere (Cantet et al., 1993; Nelsen et al., 1984,) who suggested that estimates before 1984 direct heritability were smaller than maternal heritability and genetic correlation between direct and maternal were highly negative, whereas more recent reports suggest that direct heritability estimates are larger than maternal heritability estimates with correlation between direct and maternal being also negative. Therefore the estimate in the present study are in the lower range as compared to those report in literature but agrees quite well with the notion that recent estimates of direct heritability are higher than maternal heritability estimates.

The proportion of variance due to permanent environmental maternal effects were negligible however in other species high estimates of permanent environmental maternal effects have been associated with permanent environmental effects of the uterus and the effects of multiple birth in small ruminants (Snyman et al., 1995). The estimates of ratio of maternal permanent environment variance to phenotypic variance, c^2 was of the same magnitude reported in Gobra cattle (Diop et al., 1999). Permanent environment effects results from incidence that affects all progeny of the same cow in case of birth weight (Joubert and Hammond, 1958; Ferrell 1991). The negative correlation of direct and maternal genetic effects in the present study could be explained from evolutionary point of view that it prevents species to becoming increasingly larger (Cundiff, 1972). However several workers mentioned that a possible existence of a negative environmental covariance between dam and offspring could result in a biased estimation of genetic correlation between direct and maternal genetic effects (Meyer, 1992). In this population it seemed the correlation between direct and maternal genetic effects was weak and not important. The inclusion of correlation between direct and maternal genetic in Model 4 did not change the magnitude of maternal heritability as compared to Model 3. Modeling different non genetic factors has influenced the negativity of covariance between direct and maternal genetic effects in beef cattle (Robbinson, 1996a, b; Lee and Pollak, 1997; Meyer, 1997; Dodenhoff et al., 1999). The total genetic effects for birth weight in this population was high hence genetic response could be high.

4. Conclusion

Inclusion and /or exclusion of the direct and indirect maternal genetic effects in a model may influence the magnitude of direct heritability, however the permanent environmental effects of the dam was negligible in this population. A full animal model must be used for genetic evaluation of birth weight of male Afrikaner cattle

Acknowledgement

The author would like to thank all staff who has been involved in the management of conservation experimental Afrikaner herd over the years. The herd was established in 1957 and the article was written when the first author was the head of cattle breeding section at Matopos Research Station, Bulawayo, Zimbabwe from 2001 to 2006.

References

- Bertrand, J.K., Benyshek, L.L., 1987. Variance and covariance estimates for maternally influenced beef growth traits. *J. Anim. Sci.* 64, 728.
- Brown, C.J., Galvez, M., 1969. Maternal and other effects on birth weight of beef calves. *J. Anim. Sci.* 28, 162.
- Bruckner, C.M., Slinger, W.D., 1986. Symmetric differences squared and analysis of variance procedures for estimating genetic and environmental variances and covariances for beef cattle weaning weight. II. Estimates from data set. *J. Anim. Sci.* 63, 1794-1803.
- Burfering, P.J., Kress, D.D., Friedrich, R.L., 1981. Calving ease and growth rate of Simmental-sired calves. III. Direct and maternal effects. *J. Anim. Sci.* 53, 1210.

- Cantet, R.J.C., 1990. Estimation and prediction problems in mixed linear models for maternal genetic effects. PhD. Thesis. University of Illinois, Urbana, Illinois.
- Cantet, R.J.C., Gionola, D., Misztal, I., Fernando, R.I., 1993. Estimates of dispersion parameters and genetic and environmental trends for weaning weight in Angus cattle using a maternal animal model with genetic grouping. *Livest. Prod. Sci.* 34, 203- 212.
- Cantet, R.J.C., Kress, D.D., Anderson, D.C., Doornbos, D.E., Burfening P.J., Blacwell, R.L., 1988. Direct and maternal variances and covariances and maternal phenotypic effects on preweaning growth of beef cattle. *J. Anim. Sci.* 66, 648.
- Cundiff, L.V., 1972. The role of maternal effects in animal breeding. VIII. Comparative aspects of maternal effects. *J. Anim. Sci.*, 35, 1335- 1337.
- Day, K.A., Maclaurin, G., Dube, S., Hlatshwayo, A., Trevor, C., 2003. Capturing the benefits of seasonal climate forecasts in agricultural management. Final Report for Australian Centre for International Agricultural Research (ACIAR). June 2003. Sub Project 3- Grazing Systems in Zimbabwe, 67 pp.
- Diop, M., 1997. Design and analysis of open nucleus breeding systems for cattle in Senegal. PhD Thesis, University of Nebraska, Lincoln, USA
- Dodenhoff, J., Van Vleck, L.D., Gregory, K.E., 1999. Estimation of direct, maternal and grandmaternal genetic effects for weaning weight in several breeds of cattle. *J. Anim. Sci.* 77, 840- 845.
- Gambiza, J., Nyama, C., 2000. Country pasture/forage resource profiles. Country profiles, Zimbabwe. Food and Agriculture Organization of the United Nations.
- Gestmayr, S., 1992. Impact of data structure on reliability of the estimated genetic parameters in an animal model with maternal effects. *J. Anim. Breed. Genet.* 109, 321- 336.
- Gilmour, A., 1995 Average Information Restricted Maximum Likelihood (AIREML) manual. Ploycopy.
- Groeneveld, B.E., Mostert, E., Rust, T., 1998. The covariance structure of growth traits in Afrikaner beef population. *Liv. Prod. Sci.* 55, 99- 107.
- Hagreveas, S.K., Bruce, D., Beffa, L.M., 2004. Disaster mitigation options for livestock production in communal farming systems in Zimbabwe. 1. Background information and literature review. P.O. Box 776, Bulawayo, Zimbabwe: ICRISAT and FAO Rome, Italy 56pp.
- Hlatshwayo, A., 2007. The effects of soil type and woody cover on grass production on rangelands in Zimbabwe. BSc Thesis Zimbabwe Open University, Bulawayo, Zimbabwe.
- Homann, S., Van Rooyen, A.F., Moyo, T., Nengomahsa, Z., 2007. Goat production and marketing: Baseline information for semi-arid Zimbabwe. P.O. Box 776. Bulawayo, Zimbabwe: International Crops Research Institute for the Semi-Arid Tropics. 84pp.
- Koch, R.M., Cundiff, L.V., Grgory, K.E., Dickerson, G.E., 1973. Genetic and phenotypic relations associated with preweaning and post weaning growth of Hereford bulls and heifers. *J. Anim. Sci.* 36, 235- 239.
- Lee, C., Pollack, E.J., 1997. Relationship between sire \times year interactions and direct-maternal genetic correlation for weaning weight of Simmental cattle. *J. Anim. Sci.* 75, 68- 75.
- Meyer, K., 1992a. Variance components due to direct and maternal effects for growth traits of Australian beef cattle. *Liv. Prod. Sci.* 31, 179- 204.
- Meyer, K., 1992b. Bias and sampling covariances of estimates of variance components due to maternal effects. *Genetique, Selection, Evolution.* 24, 487- 509.
- Meyer, K., 1997. Estimates of genetic parameters for weaning weight of beef cattle accounting for direct-maternal environmental covariances. *Liv. Prod. Sci.* 52, 187- 199.
- Meyer K, Garrick MJ, Donnely BJP (1994). Genetic parameters for milk production of Australian beef cows and weaning weight of their calves. *J. Anim. Sci.* 72:1155- 1165.
- Robinson, D.L., 1996b. Models which might explain negative correlations between direct and maternal genetic effects. *Liv. Prod. Sci.* 45, 111- 122.
- Robnison, D.L., 1996a. Estimation and interpretation of direct and maternal genetic parameters for weights of Australian Angus cattle. *Liv. Prod. Sci.* 45, 1-11.
- Willham, R.L., 1972. Role of maternal effects in animal breeding: III. Biometrical aspects of maternal effects in animals. *J. Anim. Sci.* 35,1288- 1293.
- Willham, R.L., 1980. Problems in estimating maternal effects. *Liv. Prod. Sci.* 7, 405- 418.