

Contents lists available at Sjournals

Scientific Journal of
Biological Sciences

Journal homepage: www.Sjournals.com



Original article

Haematologic and Biochemical Indices of *Plasmodium falciparum* Infected Inhabitants of Owerri, Imo State, Nigeria

J.O. Akaninwor^a, E.B. Essien^b, P.C. Chikezie^{b,*}, R.T. Okpara^a

^aDepartment of Biochemistry, University of Port-Harcourt, Port-Harcourt, Nigeria.

^bDepartment of Biochemistry, Imo State University, Owerri, Nigeria.

*Corresponding author; Department of Biochemistry, Imo State University, Owerri, Nigeria.

ARTICLE INFO

Article history:

Received 20 July 2013

Accepted 03 August 2013

Available online 27 August 2013

Keywords:

Haemoglobin

Packed cell volume

Erythrocyte sedimentation rate

Fasting blood sugar

Malaria

Plasmodium falciparum

ABSTRACT

The present study seeks to investigate changes in haematologic and biochemical indices of moderately *P. falciparum* infected male inhabitants of Owerri Municipality. Haematologic and biochemical indices were estimated by spectrophotometric methods. Haemoglobin concentrations of malarious subjects within age brackets of 11-20 and 21-31 years were below reference interval; [Hb]M; 11-21 years = 10.53±0.23 g/dL ($p > 0.05$); [Hb] M; 21-31 years = 11.51±1.10 g/dL ($p > 0.05$). There was no significant difference ($p < 0.05$) in erythrocyte sedimentation rate (ESR) between the two malarious groups; ESRM; 11-20 years = 29.80±0.74 mm/h; ESRM; 21-31 years = 26.51±1.42 mm/h. Packed cell volume (PCV) of malarious subject gave the following values: PCV%M; 11-20 years = 26.82±0.78; PCV%M; 21-31 years = 25.82±0.78; $p < 0.05$. Serum white blood cell count (WBC) was raised in malarious subjects compared to control groups ($p < 0.05$) except with WBC×103M; 21-30 years = 13.77±3.95; $p < 0.05$. Serum albumin was lower in malarious subjects; [Albumin]M; 11-20 years = 4.70±0.05 mg/dL and [Albumin]M; 21-31 years = 4.31±0.09 mg/dL; $p < 0.05$, whereas, serum creatinine concentrations of malarious subjects gave higher values: [Creatinine]M; 11-20 years = 0.88±0.71 mg/dL and [Creatinine]M; 21-31 years = 1.14±0.42 mg/dL; $p < 0.05$. Serum urea concentrations of malarious subjects were significantly ($p > 0.05$)

higher than the corresponding non-malarious age group. Serum fasting blood sugar (FBS) was significantly ($p > 0.05$) lower in malarious groups compared to corresponding non-malarious subjects. Specifically, [FBS]M; 11-20 years = 63.34 ± 1.66 mg/dL and [FBS]M; 21-31 years = 69.45 ± 1.25 mg/dL; $p > 0.05$. Subjects with moderate malaria infection showed symptoms of anaemia, alterations in nitrogen and carbohydrate metabolism, exemplified by raised serum level of urea and low level of FBS.

© 2013 Sjournals. All rights reserved.

1. Introduction

Four species of intracellular protozoa of the genus *Plasmodium* cause malaria in humans. They include *Plasmodium falciparum*, *P. vivax*, *P. ovale*, and *P. malariae* (Krotoski et al., 1982; Joseph et al., 2011). *P. falciparum* and *P. vivax* cause the most serious forms of the disease (WHO, 2005; Idonije et al., 2011; Joseph et al., 2011). Sporozoites from bite of female mosquitoes (genus *Anopheles*) infect humans and are the progenitor of the disease condition. The parasites have a complicated life cycle that requires a vertebrate host for the asexual cycle and female *Anopheles* mosquitoes for completion of the sexual cycle. Malaria poses a threat to public health with 80-90% of morbidity and mortality occurring in Africa, afflicting both young and old (Afolabi, 2001; Ikekpeazu et al., 2010; Ogbodo et al., 2010). In addition, reports showed that malaria could be transmitted by transfusion of infected blood (Strickland, 1991; Ali and Kadaru, 2005), sharing needles (Tracy and Webster, 2001) and congenital transmission (Ezechukwu et al., 2004).

Blood is a tissue that circulates in a virtually closed system of blood vessels. It is composed of solid elements—red, white blood cells, and platelets, suspended liquid medium—plasma. Therefore, the plasma is an extracellular fluid confined within the vascular system. The water and electrolyte composition of plasma is particularly the same as that of intracellular fluid, made up of water, electrolytes, metabolites, nutrients, proteins and hormones.

Physicochemical properties of the blood are constant but may undergo slight variations under normal physiologic conditions. However, the relative constancy in the internal environment of the blood system exhibits wide and profound perturbation and distortions under clinically defined pathophysiologic states. Some of these conditions include malignancy, genetic defects, malnutrition, parasitic infections etc. Studies have revealed that haematologic and biochemical changes occur in malaria infected blood and there are common complications associated with this disease. Haematologic changes that are associated with malaria infection include anaemia, thrombocytopenia, and disseminated intravascular coagulation (Facer, 1994; Perrin et al., 1982; Maina et al., 2010; Chandra and Chandra, 2013). Changes in physicochemical parameters of *P. falciparum* infested blood may vary with level of malaria endemicity, presence of haemoglobinopathies, nutritional status, demographic factors and level of malaria immunity (Price et al., 2001; Erhart et al., 2004). Therefore, well-informed changes in blood parameters in malaria infection enable the clinician to establish reliable diagnosis and therapeutic interventions.

Although haematologic and biochemical indices of *P. falciparum* infected individuals of Nigerian origin have been widely reported (Udesen, 2003; Egwunyenga et al., 2004; Adesina et al., 2009; Kayode et al., 2011) specific records of infected inhabitants of Owerri Municipality have been poorly documented and not widely reported in this regard. Therefore, the present study seeks to investigate changes in haematologic and biochemical indices of moderately *P. falciparum* infected male inhabitants of Owerri Municipality.

2. Materials and methods

2.1. Study area

The study was conducted between May 2011 and August 2011 in Owerri Municipality, Imo State, Nigeria, which lies on rainforest belt (Latitude 5.485° N and Longitude 7.035° E). The wet season is within the period of March-September, when breeding of *Anopheles* mosquitoes is at its peak and bites are prevalent. Twenty-one (21) clinically confirmed (WHO, 2008) and randomly selected malarious and fasting male patients attending clinics at

the Federal Medical Center (FMC), St. John Clinic/Medical Diagnostic Laboratories, Avigram Medical Diagnostic Laboratories, and Qualitech Medical Diagnostic Laboratories enrolled for this study. These centers are located in Owerri, Imo State, Nigeria. Age matched asymptomatic/non-malarious fasting male subjects (n = 15) constituted the control subjects. The patients were in the following categories- adults (n = 11) of 21-31 years old and adolescent (n = 10) of 11-20 years old. Exclusion criteria include; gastrointestinal tract infection, protein energy malnutrition, renal diseases, cirrhosis, hepatitis, obstructive jaundice, cancer, diabetes mellitus, hypertension, obesity, smoking, alcoholism, persons living with HIV, patients taking anti-malaria drugs and vitamin supplements, patients who have treated malaria in the past 2 months (Onyesom and Onyemakonor, 2011; Idonije et al., 2011) and patients with low or high parasitaemia.

2.2. Ethic

The Ethical Committee of University of Port Harcourt, Port Harcourt, Nigeria, approved the study in compliance with the Declaration on the Right of the Patient (WMA, 2000). Before enrolment for the study, the patients/subjects involved signed an informed consent form.

2.3. Collection and preparation of blood specimen

Blood specimen was collected by venipuncture from fasting subjects using 5.0 mL capacity disposable syringes. Three milliliter (3.0 mL) of the blood samples were transferred into plain bottles to allow for coagulation, whereas the remaining 2.0 mL was transferred into EDTA bottles for malaria parasite tests and haematological studies. The coagulated blood samples were centrifuged at 3000 rpm for 10 min, the serum transferred into Bijou bottle and stored frozen until required for biochemical analyses (Onyesom et al., 2010).

2.4. Malaria parasite density test

Measurement of parasite density of peripheral blood smear was by Giemsa stained techniques. The films were examined microscopically using $\times 100$ objective under oil immersion (Cheesbrough, 1998) as reported by Sumbele et al., (2010). Level of parasitaemia was in microliter (μL) of blood thick film preparation (Erhart et al., 2004). According to WHO, (2005) level of parasitaemia was graded as low+ (1 to 999 / μL), moderate++ (1000 to 9999 / μL) and severe+++ ($> 10,000$ / μL).

2.5. Haematological studies

The modified method (Baure, 1980), based on cyanomethaemoglobin reaction was used for the determination of haemoglobin concentration (Chikezie, 2009). Packed Cell Volume (PCV) was measured using whole blood mixed in a 10- μL mark capillary pipette. The set up was centrifuged at 3000 rpm for 30 min. The hematocrit was removed from the centrifuge and the volume of packed red cell column was read off and expressed as percentage of whole blood volume. Estimation of white blood cell count (WBC) was according to methods of NCCLS, (1993). Estimation of erythrocyte sedimentation rate (ESR) was according to the Westergreen's methods as described by Supcharoen et al., (1992).

2.6. Biochemical studies

Fasting blood sugar (FBS) was measured by standard methods as reported by Kazmierczack (1996). Serum urea level was determined according to the method described by Fawcett and Scott (1960) and reported by Kayode et al., (2011). Creatinine level in the blood was determined according to the methods described by Bartels et al. (1972). Albumin concentration in the blood was measured by the method of Doumas et al., (1971) and as described by Cheung and Hchman, (1996).

2.7. Statistical analyses

The experiments were designed in a completely randomized method and data collected were analyzed by the analysis of variance procedure while treatment means were separated by the least significance difference (LSD) incorporated in the statistical analysis system (SAS) package of 9.1 version (2006). The correlation coefficients between the results were determined with Microsoft Office Excel, 2010 version.

3. Results

Table 1 showed that haematological indices of non-malarious subjects were within reference intervals and there was no significant difference ($p < 0.05$) between the age brackets of 11-20 years and 21-31 years. Haemoglobin concentrations of malarious subjects within age brackets of 11-20 and 21-31 years were below reference interval; [Hb]M; 11-21 years = 10.53 ± 0.23 g/dL ($p > 0.05$); [Hb] M; 21-31 years = 11.51 ± 1.10 g/dL ($p > 0.05$). These values represented 22.97% and 31.08% drop in serum haemoglobin concentrations compared to corresponding non-malarious subjects. Serum haemoglobin concentrations between the two malarious groups were not significantly different ($p < 0.05$).

Table 1

Some haematological indices of non-malarious and malarious subjects.

Parameters	NM		M		Reference Intervals
	11-20 yrs.	21-31 yrs.	11-20 yrs.	21-31 yrs.	
[Hb] g/dL	15.67 ± 0.20^a	$16.70 \pm 0.96^{a,b}$	10.53 ± 0.23^c	$11.51 \pm 1.10^{c,d}$	13.5-18.0*
ESR mm/h	16.30 ± 1.08^a	$15.2 \pm 0.60^{a,b}$	29.80 ± 0.74^c	$26.51 \pm 1.42^{c,d}$	0-15†
PCV %	33.94 ± 0.64^a	$33.94 \pm 0.61^{a,b}$	26.82 ± 0.78^c	25.82 ± 0.78^d	40-54
WBC $\times 10^3$	6.39 ± 6.98^a	$7.53 \pm 2.26^{a,b}$	$10.13 \pm 4.75^{a,b,c}$	$13.77 \pm 3.95^{a,b,c,d}$	4.5-11.0‡

*Richards et al., (1998); †Erhart et al., (2004); ‡Bottiger and Svedberg, (1967); Means in the row with the same letter are not significantly different at $p < 0.05$ according to LSD. NM: Non-malarious; M: Malarious. WBC: cell/ μ L/mm³.

ESR of malarious subjects were above the reference intervals of ESR of ER disease as a non-specific test for acute illness and may reflect the acute process = 0-15 mm/h (Table 1) and was significantly different ($p > 0.05$) compared to the control subjects. However, there was no significant difference ($p < 0.05$) in ESR between the two malarious groups; ESRM; 11-20 years = 29.80 ± 0.74 mm/h; ESRM; 21-31 years = 26.51 ± 1.42 mm/h. PCV of malarious subject gave the following values: PCV%M; 11-20 years = 26.82 ± 0.78 ; PCV%M; 21-31 years = 25.82 ± 0.78 ; $p < 0.05$, with values below the reference interval: PCV% = 40-54. Serum WBC was raised in malarious subjects compared to control groups ($p < 0.05$) and within reference interval (WBC $\times 10^3$ = 4.5-11.0), except with WBC $\times 10^3$ M; 21-30 years = 13.77 ± 3.95 ; $p < 0.05$.

Table 2

Some biochemical indices of non-malarious and malarious subjects.

Parameters (mg/dL)	NM		M		Reference Intervals§
	11-20 yrs.	21-31 yrs.	11-20 yrs.	21-31 yrs.	
Albumin $\times 10^3$	5.18 ± 0.29^a	$4.46 \pm 0.05^{a,b}$	$4.70 \pm 0.05^{a,b,c}$	$4.31 \pm 0.09^{a,b,c,d}$	3.5-5.5
Creatinine	0.62 ± 0.27^a	$0.94 \pm 0.51^{a,b}$	$0.88 \pm 0.71^{a,b,c}$	$1.14 \pm 0.42^{a,b,c,d}$	0.7-1.5
Urea	10.70 ± 0.94^a	$12.72 \pm 0.51^{a,b}$	17.10 ± 0.74^c	26.14 ± 0.98^d	8-20
FBS	89.42 ± 0.64^a	$87.47 \pm 1.06^{a,b}$	63.34 ± 1.66^c	69.45 ± 1.25^d	60-100

§Martin, (1983): Means in the row with the same letter are not significantly different at $p < 0.05$ according to LSD. NM: Non.

Table 2 showed that there was no significant difference ($p < 0.05$) between the two non-malarious groups in connection to the four experimental biochemical indices. Likewise, serum albumin and creatinine concentrations were not significantly different ($p < 0.05$) between the non-malarious and malarious subjects. In addition, marginal changes in serum albumin and creatinine concentrations in non-malarious and malarious subjects were within reference intervals (Table 2). Specifically, serum albumin was lower in malarious subjects; [Albumin]M; 11-20 years = 4.70 ± 0.05 mg/dL and [Albumin]M; 21-31 years = 4.31 ± 0.09 mg/dL; $p < 0.05$, whereas, serum creatinine concentrations of malarious subjects gave higher values: [Creatinine]M; 11-20 years = 0.88 ± 0.71 mg/dL and [Creatinine]M; 21-31 years = 1.14 ± 0.42 mg/dL; $p < 0.05$. Serum urea concentrations of malarious subjects were significantly higher than the corresponding non-malarious age group. Serum urea concentration of malarious subjects between the age brackets of 21-31 years was above the reference interval; [Urea]M; 21-31 years = 26.14 ± 0.98 mg/dL; $p > 0.05$. Serum FBS concentration was significantly ($p > 0.05$) lower in malarious groups compared to corresponding non-malarious subjects. Specifically, [FBS]M; 11-20 years = 63.34 ± 1.66 mg/dL and

[FBS]M; 21-31 years = 69.45±1.25 mg/dL; $p > 0.05$. These values represented 29.17% and 20.60% drop in serum FBS concentrations compared to their corresponding non-malarious age group.

4. Discussion

Haematologic changes associated with malaria infection are well recognized and have been widely reported (Das et al., 1999; Mishra et al., 2002; Udosen, 2003; Bidaki and Dalimi, 2003; Erhart et al., 2004; Maina et al., 2010). The present study reported haematologic and biochemical changes associated with moderate malaria infection in male subjects. The decreased haemoglobin concentrations in malarious subjects (Table 1) were predictive as had been reported by several authors (Das et al., 1999; Mishra et al., 2002; Udosen, 2003; Bidaki and Dalimi, 2003; Erhart et al., 2004; Maina et al., 2010). Earlier reports had posited that malaria-related anaemia is often more severe in areas of intense malaria transmission and affects younger children rather than older children or adults (Phillips and Pasvol, 1992; Menendez et al., 2000). Also, the present study showed that moderate malaria infection among male subjects in Owerri Municipality exhibited the same pattern as described above. Moderate infection caused reduction in haemoglobin concentration, which was more pronounced in adolescents between the age brackets of 11-20 years than their adult counterparts of age brackets between 21-30 years ($p < 0.05$) (Table 1). According to reported by Maina et al., (2010) as contained in the National Guidelines for Diagnosis, Treatment and Prevention of Malaria For Health Workers in Kenya, anaemia is defined as [Hb] < 10 g/dL for both males and females. Furthermore, severe malaria anaemia is defined as [Hb] < 5 g/dL in the presence of hyperparasitaemia ($> 200,000$ parasites/ μ L). Therefore, the drop in haemoglobin concentrations in the malarious subjects (Table 1) approximately connoted mild anaemia. The decreased PCV levels were also expected from previous reports (Adesina et al., 2009; Ogbodo et al. 2010; Kayode et al., 2011). Furthermore, the drop in PCV values in the two malarious groups confirmed symptoms of anaemia in these study groups. Two striking factors are responsible for the development and presentation of anaemia in malaria infections.

Rapid rate of haemolysis associated with the pathophysiology of the disease condition (Phillips and Pasvol, 1992; Selvam and Baskaram, 1996; Erhart et al., 2004).

Reduced rate of haemoglobin biosynthesis, which is often connected to level of immunity and nutritional status of infected individuals (Das et al., 1999; Price et al., 2001; Wickramasinghe and Abdalla, 2000; Erhart et al., 2004).

Therefore, the interplay of these multifactorial etiologies of anaemia in malaria infection, as described above, was responsible for the drop in haemoglobin concentrations in the malarious groups by 22.97% and 31.08% (Table 1). In concord with the present findings, studies among non-immune or semi-immune populations outside Africa have also shown statistically significant levels of mild anaemia in falciparum malaria patients (Rojanasthien et al., 1992; Das et al., 1999).

Elevation of ESR have been reported in acute and chronic infections (Kwiatkoski et al., 1989), chronic inflammatory disorders (Kwiatkoski et al., 1989; Supcharoen et al., 1992; Dreyer and Boden, 2003) malignancies especially Hodgkin's disease (Malcolm and Brigden, 1999; Mönig et al., 2002; Dreyer and Boden, 2003), tissue necrosis (Scuderi, 1986; Beutler and Cerami, 1987) and pregnancy (van den Broek and Letsky, 2008). Supcharoen et al., (1992), used ESR as basis for the diagnosis and monitor of therapeutic intervention of malaria. They suggested that ESR was elevated during acute malaria infection and declined with recovery. Thus, the present findings as presented in Table 1 was in agreement with the reports of Supcharoen et al., (1992). However, measurement of ESR is often used as a non-specific test for acute illness and may reflect the acute process of the disease.

Erhart et al., (2004) stated that semi-immune persons in Western Thailand with parasitaemia tended to have significantly lower white blood cell. Perrin et al., (1982) and Rojanasthien et al., (1992) reported contrary findings during malaria infection in man. The non-significant ($p < 0.05$) increase in serum WBC in the present study contradicts these two separate reports. Nevertheless, the present study showed that the malarious subjects did not exhibit leukocytosis, which was defined as total WBC $> 17,000/\mu$ L, frequently seen in 8% malarious individuals as against 3% non-malarious children living in Western Kenya (Maina et al., 2010). In another study, Kayode et al., (2011) indicated significant increase ($p < 0.05$) in WBC of malaria and malaria typhoid co-infected patients, which they posited could have been elicited by increased production of leukocytes at the onset of the infection to wade off malaria parasite and typhoid pathogens. Similarly, increase in WBC in pregnant and non-pregnant malaria patients has been reported by Adesina et al., (2009) and Sumbele et al. (2010). However, the works of Ali et al.

(2009), noted both increased and decreased WBC in the blood of typhoid patients examined in Dubai. From these indications, the use of serum level of WBC as an index for diagnosis may not be very reliable. Therefore, WBC should always be thoroughly re-evaluated for malaria for reproducibility and reliability.

Studies carried out by Amah et al., (2011) showed significant reduction in serum levels of albumin in malaria patients in endemic regions of Calabar, Nigeria. Many authors have proposed the use of serum albumin levels as reliable biochemical marker for establishing severe pathologic conditions such as malnutrition and infectious diseases (Das et al., 1997; Kwena et al., 2012). Malaria infections are accompanied with significant decrease in plasma albumin concentrations (Kwena et al., 2012) as well as in malnutrition and pregnancy. However the prevailing plasma albumin concentration in malaria infection is dependent on the nutritional status of the affected individual and hepatic functionality (Crawly, 2004; Ogbodo et al., 2010). Probably, based on the nutritional and hepatic status of the experimental subjects, the report presented here showed non-significant ($p < 0.05$) reduction in serum albumin levels in malarious subjects compared to non-malarious groups (Table 2). Contrary to these observations, Ogbodo et al., (2010) showed that there was initial significant ($p > 0.05$) increase in serum levels of albumin in low and moderate malaria infections, but decreased as the malaria density increased. Based on these observations, they recommended the use of albumin infusion in place of other colloidal solutions as a good intervention in severe malaria.

Creatinine and urea are nitrogenous low threshold substances with immense clinical application in ascertaining renal function. Impairment of renal function during severe falciparum malaria is common (al-Yaman et al., 1997; Eiam-Ong and Sitprija, 1988; Günther et al., 2002; Mockenhaupt et al., 2004). Table 2 shows that the malarious groups presented marginal increases in serum creatinine concentration ($p < 0.05$). Paradoxically, serum levels of urea was significantly ($p > 0.05$) raised in the same malarious subjects under investigation. But elevation of serum urea concentration could also connote evidence of dehydration, consumption of proteinous meals and tissue catabolism. Nevertheless, this was an obvious indication that moderately malaria infected subjects exhibited alteration in nitrogen metabolism with underlying compromised renal function. According to Sitprija, (1988), raised blood urea concentration reflected gradual progression towards renal dysfunction. Specifically, serum levels of urea had been observed to increase more rapidly than serum creatinine concentration in individuals with renal dysfunction (Emian-Ong, 2002).

Blood sugar levels in malaria infection has received the attention of several researchers. Studies by Kayode et al., (2011) indicated hypoglycemia in both malaria and typhoid co-infected patients. They posited that the level of hypoglycemia correlated with severity of infection, which was elicited by hyper-secretion of insulin. Their report corroborates the studies by Onyesom and Agho (2011), who noted the incidence of hypoglycemia in malaria patients in Edo-Delta state. The role of low serum insulin-like growth factor-1 (IGF-1) and low blood glucose levels in malaria infection was reported by Mizushima et al., (1994). They noted that *P. falciparum* infected children with low IGF-1 levels (< 50 ng/mL) presented hypoglycemia compared to other study groups. In another study, Binh et al., (1997) reported the relative contribution of insulin-mediated and non-insulin-mediated plasma glucose levels in severe malaria. The report stated that there was a corresponding increase risk of hypoglycemia as infection progressed because of host glucose production becomes insufficient for host/parasite demand. The study also revealed that basal plasma glucose increased in uncomplicated malaria, because of peripheral insulin resistance. Moderate malaria infection caused significant reduction in serum FBS levels (Table 2) with blood sugar levels tending towards hypoglycemia ([FBS] < 60 mg/dL). Accordingly, the present report supports the findings of previous authors (Mizushima et al., 1994; Binh et al., 1997; Kayode et al., 2011; Onyesom and Agho 2011).

Finally, although these changes in haematologic and biochemical indices in association with malaria infection are not novel, our findings have added more information, hitherto the limited knowledge and sparsely reports on changes in blood profile of malaria infected individuals habitat in Owerri Municipality .

References

- Adesina, K.T., Balogun, O.R., Babatunde, A.S., Sanni, M.A., Fadeyi A., Aderibigbe, S., (2009). Impact of malaria parasitaemia on haematologic parameters in pregnant women at booking in Ilorin, Nigeria. *Trends in Medical Research*. 4, 84-90.
- Afolabi, B.M. (2001). Malaria: The global scourge. *Nigeria Clinic*. 5, 9-12.

- Ali, H.A., Ahmed, M.S.A. Jawahar, L.G. Abdulla, M.U., Nadeem, J.Y. and Hina, S.H. 2009. Hematological and biochemical changes in typhoid fever. *Pakistan. J. Med. Sci.* 25, 166-171.
- Ali, M.S.M., Kadaru, A.G.M. 2005, in vitro processing of donor blood with sulphadoxine/pyrimethamine for eradication of transfusion induced malaria. *America J.Trop. Med. Hyg.* 73(6), 1119 – 1123.
- al-Yaman, F., Awburn, M.M., Clark, I.A. 1997. Serum creatinine levels, reactive nitrogen intermediates in children with cerebral malaria in Papua New Guinea. *Trans. Royal. Soc.Trop. Med Hyg.* 91(3), 303-305.
- Amah, U.K., Ahaneku, J.E., Usoro, C.A.O., Ezeoke, A.C.J., Okwara, J.E., Amah, A.K., Etukudo, M.H., Okwara, E.C., Amah, B.C. Comparative study of C-reactive protein and other biochemical parameters in patients with hepatitis B and malaria in Calabar, Nigeria. *Nigeria J. Physiol. Sci.* 26, 109 – 112.
- Bartels, H., Bohmer, M., Heierli, C. 1972. Serum creatinine determination without protein precipitation. *Clinica. Clin. Chim. Acta. Acta*, 37, 193-197.
- Baure, J.D. 1980. Laboratory investigation of hemoglobin. In: *Gradwohl's Clinical Laboratory Methods*, Diagnosis, Editors. Sonnenwirth, A.C., Jarett, L. St. Louis, MO: Mosby.
- Beutler, B. and Cerami, A. 1987. Cachectio: More than a tumor necrosis factor. *New. Engl. J. Med.* 379-85.
- Bidaki, Z.M., Dalimi, A.A. 2003. Biochemical, hematological alteration in Vivax malaria in Kahnouj city. *J.Rafsanjan. U. Med. Sci.* 3, 17-24.
- Binh, T.Q., Davis, T.M., Johnston, W., Thu, L.T., Boston, R., Danh, P.T., Anh, T.K. 1997. Glucose metabolism in severe malaria: minimal model analysis of the intravenous glucose tolerance test incorporating a stable glucose label. *Metabol.* 46(12), 1435-40.
- Bottiger, L.E., Svedberg, C.A. 1967. Normal erythrocyte sedimentation rate and age. *Br. Med. J.* 2, 85–87.
- Chandra, S., Chandra, H. 2013. Role of haematological parameters as an indicator of acute malarial infection in Uttarakhand State of India. *Mediterr. J.Hematol. Infect. Dis.* 5(1), e2013009, DOI 10.4084/MJHID.2013.009.
- Cheesbrough, M., 1998. *District laboratory practice in tropical countries*. Cambridge University Press, Cambridge.
- Cheung, K. and Hchman, P.E. 1996. *Methods of Analysis of Albumin: Liver Function*. In: *Clinical chemistry, theory, analysis and correlation*. Kaplan, L.A., Pesce, A.J. (Eds.). 3rd Edn, Mosby, London.
- Chikezie, P.C. (2009). Comparative methaemoglobin concentrations of three erythrocyte genotypes .HbAA, HbAS and HbSS, of male participants administered with five antimalarial drugs. *African. J. Biochem. Res.* 3(6), 266-271.
- Crawly J. 2004. Reducing the burden of malaria in infants, young children in malaria endemic countries of Africa: from evidence to action. *America. J. Trop. Med.Hyg.* 71 (2 Suppl), 25 -34.
- Das, B.S., Nanda, N.K., Rath, P.K., Satapathy, R.N., Das, D.B. 1999. Anaemia in acute, Plasmodium falciparum malaria in children from Orissa state, India. *Annals of Tropical Medicine and Parasitology.* 93, 109–119.
- Das, B.S., Thurnham, D.J. and Das, D.B., 1997. Influence of malaria on markers of iron status in children: implications for interpreting iron status in malaria–endemic communities. *Br. J. Nutr.* 78, 751-760.
- Doumas, B.T., Watson, W.A. and Biggs, H.G. (1971). Albumin standard and measurement of serum albumin with bromocresol green. *Clin. Chim. Acta.* 31, 87-96.
- Dreyer, S.J. and Boden, S.D. 2003. Laboratory evaluation in neck pain. *Phy. Med.Rehabilitation. Clin. N. America.* 14, 589–604.
- Egwynyenga, A.O., Isamah, G., Nmorsi, O.P. (2004). Lipid peroxidation, ascorbic acid levels in Nigeria children with acute falciparum malaria. *Afr. J. Biotechnol.* 3, 560-563.
- Eiam-Ong, S. and Sitprija, V. 1998. Falciparum malaria and the kidney: a model of inflammation. *American. J. Kidney. Diseases.* 32, 361–375.
- Emian-Ong, S. 2002. Current knowledge of falciparum malaria-induced acute renal failure. *J. Med Assoc Thai. Suppl.* 16, 24.
- Erhart, L.M., Yingyuen, K., Chuanak, N., Buathong, N., Laoboonchai, A., Miller, R.S., Meshnick, S.R., Gasser, Jr R.A., Wongsrichanalai, C. 2004. Hematologic and clinical indices of malaria in a semi-immune population of western Thailand. *America. J. Trop. Med. Hyg.* 70(1), 8–14.
- Ezechukwu, C., Ekejindu, E., Ugochukwu, E. and Oguatu, M. (2004). Congenitally acquired malaria in hyperendemic area. A Cohort study: *Trop. J. Med. Res.* 8(2), 44-48.
- Facer, C.A. 1994. Hematological aspects of malaria. In: *Infection and Hematology*. Oxford: Butterworth Heinemann Ltd.
- Fawcett, J.K., Scott, J.E. 1960. A rapid, precise method for the determination of urea. *J. Clin. Pathol.* 13, 156-159.

- Guñther, A. Burchard, G. D. Slevogt, H. Abel, W., Grobusch, M. P. 2002. Renal dysfunction in falciparum–malaria is detected more often when assessed by serum concentration of cystatin C instead of creatinine. *Trop. Med. Inter. Health.* 7(11), 931–934.
- Idonije, O.B., Festus, O., Okhiai, O., Akpamu, U. 2011. Comparative study of the status of a biomarker of lipid peroxidation .malondialdehyde. in patients with Plasmodium falciparum and Plasmodium vivax malaria infection. *Asian. J. Biol. Sci.* 4, 506-513.
- Ikekpeazu, E.J., Neboh, E.E. Maduka, I.C. Nwagbara, I.J., Nwobodo, M.W. 2010. Type-2 diabetes mellitus and malaria parasitemia: Effect on liver function tests. *Asian. J. Med. Sci.* 2(5), 214-217.
- Joseph, V., Varma, M., Vidhyasagar, S., Mathew, A. 2011. Comparison of the clinical profile and complications of mixed malarial infections of Plasmodium falciparum and Plasmodium vivax versus Plasmodium falciparum mono-infection. *Sultan Qaboos University. Med. J.* 11(3), 377–382.
- Kayode, O.T. Kayode, A.A.A., Awonuga, O.O. 2011. Status of selected hematological and biochemical parameters in malaria and malaria-typhoid co-infection. *J. Biol. Sci.* 11, 367-373.
- Kazmierczack, S.C., 1996. Methods of analysis of creatinine, urea, total protein, inorganic phosphate, calcium and pH.: renal function. In: *Clinical Chemistry, Theory, Analysis and Correlation*, Kaplan, L.A. and A.J. Pesce, (Eds.). 3rd Edn. Mosby, London.
- Krotoski, W.A., W.E. Collins, R.S. Bray, P.C. Garnham and F.B. Cogswell et al., 1982. Demonstration of hypozoites in sporozoite transmitted Plasmodium vivax infection. *America. J. Trop. Med.Hyg.* 31, 1291-1293.
- Kwena, A., Wakhisi, J. and Mambo, F. 2012. Possible Biochemical Markers in Plasmodium falciparum Malaria Infected Children with or without Malnutrition at Webuye and Eldoret, Western Kenya. *Adv. Bioresearch.* 3(2), 49 – 54.
- Kwiatkoski, D., Cannon, J.G., Manogue, K.R, et al.1989. TNF production in falciparum malaria and its association with schizont rupture. *Clin. Exp. Immunol.* 77, 391-396.
- Maina, R.N, Walsh, D., Gaddy, C., Hongo, G., Waitumbi, J., Otieno, L., Jones, D. and Ogutu, B.R. 2010. Impact of Plasmodium falciparum infection on haematological parameters in children living in Western Kenya. *M. J.* 9, S4. <http://dx.doi.org/10.1186/1475-2875-9-S3-S4>.
- Malcolm, L. and Brigden, M.D. (1999). Clinical utility of the erythrocyte sedimentation rate. *American Family Physician.* 60(5), 1443-1450.
- Martin, D.W. 1983. Blood plasma., clotting. In: Martin, D.W., Mayes, P. A., Rodwell, V.W. editors. *Harper’s Review of Biochemistry.* 9th ed. California: Lange. Med. Publ.
- Menendez, C., Fleming, A.F., Alonso, P.L. 2000. Malaria-related anaemia. *Parasitology Today.* 16, 469–476.
- Mishra, S.K., Mohaptra, S., Mohantu, S., Patel, N.C., Mohaptra, D. N. 2002. Acute renal failure in falciparum malaria. *J. Indian. Acad. Clin. Med.* 3, 141-147.
- Mizushima, Y., Kato, H., Ohmae, H., Tanaka, T., Bobogare, A., Ishii, A. 1994. Prevalence of malaria ., its relationship to anaemia, blood glucose levels and serum somatomedin C (IGF-1) levels in the Solomon Islands. *Acta. Trop.* 58(3-4), 207-210.
- Mockenhaupt, F., Ehrhardt, S., Burkhardt, J., Bosomtve, S., Laryea, S., Anemana, S., Otchwemah, R., Cramer, J., Dietz, E., Gellert, S. and Bienzle, U. 2004. Manifestation ., outcome of severe malaria in children in Northern Ghana. *America. J. Trop. Med. Hyg.* 71(2): 167-172.
- Mönig, H., Marquardt, D., Arendt, T., Kloehn, S. 2002. Limited value of elevated erythrocyte sedimentation rate as an indicator of malignancy. *Fam. Pract.* (5), 436-438.
- National Committee for Clinical Standards. 1993. Reference procedure for erythrocyte sedimentation rate (ESR) test, 3rd ed. H2-A3. Villanova, Pa. NCCLS.
- Ogbodo, S. O., Okeke, A. C., Obu, H. A, Shu, E.N. and Chukwurah E.F. 2010. Nutritional status of parasitemic children from malaria endemic rural communities in eastern Nigeria. *Curr. Pediatr. Res.* 14(2), 131-135.
- Onyesom, I., Agho, J.E. 2011. Changes in serum glucose and triacylglycerol levels induced by the co-administration of two different types of antimalarial drugs among some malarial patients in Edo-Delta Region of Nigeria. *Asian J.Sci. Res.* 4, 78-83.
- Onyesom, I., Onyemakonor, N. 2011. Levels of parasitaemia and changes in some liver enzymes among malarial infected patients in Edo-Delta Region of Nigeria. *Curr. Res. J. Biolog. Sci* 3(2), 78-81.

- Onyesom, I., Ekeanyanwu, R.C., Achuka, N. 2010. Correlation between moderate Plasmodium falciparum malarial parasitaemia and antioxidant vitamins in serum of infected children in South Eastern Nigeria. *African J.Biochem. Res.* 4(12), 261-264.
- Perrin, L.H., Mackey, L.J., Miescher, P.A. 1982. The hematology of malaria in man. *Semin Hematology.* 19, 70–82.
- Phillips, R.E., Pasvo, G. 1992. Anaemia of Plasmodium falciparum malaria. *Baillieres. Clin. Haematol.* 5, 315–330.
- Price, R.N., Simpson, J.A., Nosten, F., Luxemburger, C., Hkirjaroen, L., Kuile, F., Chongsuphajaisiddhi, T., White, N.J. 2001. Factors contributing to anaemia after uncomplicated falciparum malaria. *America. J.Trop. Med. Hyg.* 65, 614–622.
- Richards, M.W., Behrens, R.H., Doherty, J.F. 1998. Short report: Hematologic changes in acute, imported Plasmodium falciparum malaria. *America. J. Trop. Med. Hyg.* 59, 859.
- Rojanasthien, S., Surakamollear, V., Boonpucknavig, S., Isarangkura, P. 1992. Hematological and coagulation studies in malaria. *J.Med. Assoc. Thai.* 75, 190–194.
- Scuderi, P., Sterling, K., Lamks, S. 1986. Raised serum levels of TNF in parasitic infections. *Lancet.* 2, 1364-1365.
- Selvam, R., Baskaran, G., 1996. Hematological impairments in recurrent Plasmodium vivax infected patients. *Japan. J.Med. Sci.Biology.* 49, 151–165.
- Sitprija, V., 1988. Nephrology in Falciparum malaria. *Kidney. Int.* 34, 867-877.
- Strickland, G.T. 1991. Life cycle of malaria parasite. In: *Tropical Medicine.* 7th Edn, W.B. Saunders, USA, 586-601.
- Sumbele, I.U.N., Theresa, N.A. Samje, M. Ndzeize, T. Ngwa, E.M., Titanji, V.P.K., 2010. Hematological changes and recovery associated with untreated and treated plasmodium falciparum infection in children in the mount Cameroon region. *J.Clin. Med. Res.* 2, 143-151.
- Supcharoen, O., Widjaja, H., Ali, K.B., Kitayaporn, D., Pukrittayakamee, S., Wilairatana, P Sathawarawong, W., 1992. A study of erythrocyte sedimentation rate in malaria. *J. Infect. Dis. Anti. Microbrobial. Agents.* 9(4), 193-199.
- Tracy, J.W., Webster, L.T. 2001. Drugs used in the chemotherapy of protozoan infections. In: (Adam, J.G., Limbird, L.E. and Gilman, A.G. (Eds). *Goodman and Gilman’s Pharmacological Basis of Therapeutics.* 10th Edition, McGraw-Hill, U.S.A.
- Udosen, E.O., 2003. Malaria treatment using oral Medkafin: Changes in biochemical and hematological parameters in Nigerian children with uncomplicated falciparum malaria. *Orient, J. Med.* 15, 12-22.
- van den Broek, N.R., Letsky, E.A. 2008. Pregnancy ., the erythrocyte sedimentation rate. *BJOG: An Int. J. Obstet. Gynaecol.* 108(11), 1164–1167. DOI: 10.1111/j.1471-0528.2003.00267.x.
- WHO .2005. Susceptibility of Plasmodium falciparum to antimalarial drugs: report on global monitoring, 1996–2004. *WHO/HTM/MAL/2005.1103.*
- WHO . 2008. Severe P. falciparum malaria. *Transaction. Roy. Soc. Trop. Med.Hyg.* 94, 51-59.
- Wickramasinghe, S.N., Abdalla, S.H. 2000. Blood and bone marrow changes in malaria. *Baillieres. Best. Pract. Res. Clin. Haematol.* 13, 277–299.
- WMA., 2000. World medical association declaration of Helsinki ethical principles for medical research involving human subjects. 52nd WMA General Assembly, Edinburgh, Scotland.