

**Original article****Molecular detection of *Candidatus liberibacter asiaticus* associated with Citrus greening (Huanglongbing) of mandarin by DNA template preparation****G.P. Jagtap*, M.V. Mahajan, U. Dey**

Department of Plant Pathology, M.K.V., Parbhani, Maharashtra, India.

*Corresponding author; Department of Plant Pathology, M.K.V., Parbhani, Maharashtra, India.

ARTICLE INFO

ABSTRACT

Article history:

Received 03 December 2012

Accepted 18 January 2013

Available online 31 January 2013

Keywords:

PCR

HLB

Citrus

The polymerase Chain Reaction (PCR) diagnosis is more reliable and sensitive diagnostic tool for detecting greening bacterium than other conventional approaches like Electron microscopy, DNA-DNA hybridization and immunofluorescence (IF) for detection of citrus greening. During experiments, it was observed that sodium sulphite method of DNA isolation provided higher yield and better quality DNA than other methods. Primer C (450 bp) was more efficient in amplifying the DNA of greening bacterium even at a very low concentration of 0.1 µg. To confirm the reliability of PCR, the greening bacterium was also detected in graft-inoculated plants, which showed typical greening bacterium was also detected in graft-inoculated plants, which showed typical greening symptoms. Results showed amplification of 450 bp in PCR suggesting sampling in March is more suitable for PCR detection of greening bacterium.

© 2013 Sjournals. All rights reserved.

1. Introduction

Citrus belongs to family Rutaceae and has approximately 150 genera and 1600 species, which are found in tropical, temperate regions of the world. Globally citrus is grown in 114 countries. Out of these, 53 countries grow citrus commercially. On production basis, Brazil tops the list amongst citrus producing countries with 19.9 million tons followed by USA with 14.1 million tons, China with 12.1 million tons, Mexico with 6.4 million tons, Spain with 5.7 million tons and India 4.4 million tons at the 6th position.

Citrus is the third important fruit crop of India after banana and mango with the acreage of 5.63 lakh ha and production of 56.8 lakh tons and average productivity of 10.1 t/ha. The most important commercial citrus cultivars in India are, the mandarin (*Citrus reticulata* Blanco), followed by sweet orange (*Citrus Sinensis osbeck*) and acid lime (*Citrus aurantifolia* Swingle) sharing 65, 25 and 10 per cent, respectively, of all citrus fruits produced in the country. The state of Maharashtra is considered as the top most citrus growing state in terms of area (2.38 lakh ha) and second in terms of production (14.4 lakh tones) and 10th in terms of productivity (6.0 t/ha) according to Indian Horticulture Database, 2005.

In India, the major pathogens of economic importance in citrus are citrus tristeza virus (CTV), citrus yellow mosaic badna virus (CYMV), indian citrus ring spot virus (ICRSV), Viroids diseases like citrus exocortis viroid and a fastidious prokaryote causing citrus greening disease. Citrus greening disease is an important disease of citrus which greatly affects the production of citrus fruits in several parts of India (Ahlawat, 1997). In India, the greening disease was first identified by L.R. Fraser in 1965 and its wide spread occurrence was confirmed by Varma *et al.* (1993), Ahlawat and Pant (2003).

Confirmation that a citrus tree is affected by greening has up to now relied on the electron microscopical identification of the bacterium, but due to erratic distribution of the pathogen in sieve tubes, it was cumbersome process to cut the right tissue having greening bacterium. However, several indirect approaches such as biological indexing, immunofluorescence tests with monoclonal antibodies and DNA-DNA hybridization with radioprobes have been used for greening diagnosis (Ahlawat and Pant, 2003). Biological indexing is a time consuming procedure and temperature dependent. It requires a well-equipped glass houses and long term maintenance of indicator hosts. The use of monoclonal antibodies for field diagnosis has proven unsatisfactory (Korsten *et al.*, 1993). Detection by DNA probes, though an accurate method for detection, but requires handling of radioactive elements and is being discouraged. Moreover, these are not practically feasible methods for handling a large sampling unit. The recent study by Hocquellet *et al.* (2000) and Ahlawat *et al.* (2003) have shown that gene amplification of β operon ribosomal protein is a sensitive and promising technique for detection and differentiation of greening bacterium. However more work is required for standardization of detection of greening organism Polymerase Chain Reaction (PCR) and its application on large scale.

2. Materials and methods

2.1. Confirmation of infection in trees

The three trees showing typical symptoms of greening were identified and labeled. The symptomatic leaves from individual tree were taken as source material for extraction of DNA and confirmation of infection by PCR as described by Ahlawat *et al.*, (2003).

2.2. Graft transmission

The scions from these six trees were grafted on mandarin sweet orange in the glasshouse. The observation for symptoms development in grafted plants was taken periodically and finally grafted plants were tested using PCR.

2.3. DNA extraction from citrus tissue

The total DNA was isolated from midrib and petiole of symptomatic leaves of infected field trees. Three methods of DNA extraction viz. DN-easy Plant Mini kit (QIAGEN, Germany), sodium sulphite (Baranwal *et al.*, 2003), Nucleic Acid Technique were employed.

2.4. DNA extraction protocol by commercial kit (QIAGEN DN-easy Plant Mini kit)

150 from leaves of greening infected midrib were ground in liquid nitrogen in sterilized pestle and mortar. The tissue powder was transferred into 1 ml eppendorf tube. The DNeasy membrane and it was incubated for 5 min at room temperature and centrifuge for 1 min at 6000 rpm. The above step was repeated. DNA was isolated according to Qiagen's manual.

2.5. DNA extraction by sodium sulphite method (Baranwal et al., 2003)

10 ml extraction buffer was prepared (Appendix) in a single Mortar and pestle. 10 ml extraction buffer was heated to 65°C before adding to powdered tissue. The powdered tissue was taken into 1.5 eppendorf tube. 1 ml hot extraction buffer was added. The eppendorf tube containing powdered tissue in extraction buffer was kept at 95°C heating block for 10 min. Regular vortexing after every 2 min was done and kept on ice for 2 min. The tube was centrifuged at 12000 rpm for 5 min. Approximately 800 µl of supernatant was taken and transferred to new tube containing 5 µl of RNase. Incubated at 37°C for 20 min. 480 µl of isopropanol was added to it and mixed by gentle rocking. The tube was centrifuged for 5 min at 12000 rpm. To the pellet, 30 µl of sterile distilled water was added. To dissolve, DNA pellet with distilled water was heated briefly at 50°C and flicking was done. The DNA was precipitated with 30 µl of 3M solution acetate and 1/10th volume of 95 per cent of ethanol. The tube was kept on ice for 10-20 min. The tube was centrifuged for 5 min with 12000 rpm. The ethyl alcohol was poured off. The pellet was dried for 40 min at 37°C. The pellet was dissolved in 70-100 µl of double distilled water, reheated 45-50°C.

2.6. DNA extraction using membrane bard nucleic acid technique (Baranwal, Gupta and Singh, 2007)

Take 100 mg of petiole and midrib of leaves tissue from las infected plant were homogenized in 1 ml of alkaline solution of NaOH. The resulted extract were incubated at room temperature (24-32°C) for 15 min or centrifuged at 1200 g for 10 min. 5 µl of sap were spotted on untreated NCM5 (BAS 85, poresize 0.45 µm Scwicher and Schuee, Kece, N.H.) that were dried for 30 min at 24-32°C. Individual spot (4.0 mm) for each sample were cut out with paper hole punch (Kangaro industries, Ludhiana, India) and eluted in 30 µl of sterile distilled water by incubation at 80°C for 10 min on a heat block. The liquid was collected by centrifugation (termed NCM eluted extract). Volumes of 2.5, 5, 10 and 20 µl were used for detection of clA ANAs by PCR.

2.7. Primer synthesis

Pair of primers from conserved region of ribosomal β-operon gene and ribosomal DNA was synthesized and used to detect citrus greening bacterium in PCR system (CG3450F) primer (Table 1).

2.8. Ploymerase enzyme

Polymerase enzyme (Taq polymerase) was evaluated for their efficiency to amplify and it was used with primer to compare their efficacy.

2.9. PCR amplification

The amplification was performed on a thermal cycle (manufacture) using primer and polymerase enzyme (Taq) for comparing their efficiency in amplification of DNA of greening bacterium based on the number of sample amplified and intensity of amplified DNA band.

The conditions followed for PCR are as follows. Initial denaturation was given at 94°C for 5 minutes 1 cycle followed by denaturation at 94°C for 30 seconds, annealing at 60°C for 30 seconds and extension of 72°C for 1 minute and the final extension was given at 72°C for 10 minutes 30 cycles.

Table 1

Specific primers used for amplification of ribosomal β-operon gene and 16s ribosomal DNA of citrus greening bacterium

Primers	Primer sequence	Annealing temperature	Amplicon size
A	5'GCGCGTATCCAATACGAGCGGCA3' 5'GCCTCGCGACTTCGCAACCCAT3'	62°C	1160bp
B	5'TATAAAGGTTGACCTTTCGAGTTT3' 5'ACAAAAGCAGAAATAGCAAACA3'	58°C	703bp
C	5'TGGGTGGTTTACCATTCAAGT3' 5'CGCGACTTTCGCAACCCATTG'	58°C	450bp

2.10. Analysis of PCR product by electrophoresis

Following PCR, amplicon were analyzed in 1 % agarose gel and electrophoresed in Tris-acetate EDTA (TAE) buffer containing ethidium bromide 0.5 g agarose was melted in 50 ml 1x TAE running buffer and 2µl ethidium bromide was added to it after cooling to around 50^oC and poured into a casting tray for polymerization placing the 12 well combs. The comb was removed after polymerization and the gel was then placed on electrophoresis tray filled with 1 x PAE buffer 20 µl each of PCR product mixed with 2 µl of 6 x loading dye was loaded into the well and was run at 60 volt for 30 min. A aliquot of 1 kb DNA ladder 4 µl was named with dye similarly and electrophoresed to serve as molecular weight marker. After the run the gel was observed under ultraviolet (UV) transillumina and photographed on thermal paper using gel documentation system.

2.11. Validation of PCR for detection of greening bacterium

Following the standardization of DNA extract method primer and enzyme was validated by taking 40 random sample collections from the apparently infected mandarin trees in orchard.

3. Results and discussion

3.1. Confirmation of the presence of greening in infected trees

The trees were identified during survey and tested for the presence of greening inoculum by PCR. It was found that the pathogen was found from all the trees. The details of the PCR will be given in subsequent results. The leaves from these trees were collected and used in further experiments.

3.2. Graft transmission

Of the Six-PCR positive trees, grafted on mandarin plants showed typical greening symptoms in 3-6 months after grafting. The new leaves showed yellowing followed by mottling of leaves. of the grafted plant, plants showed amplification of greening pathogen in PCR when tested after 6 months. However, since the studies were planned with a view to develop PCR diagnostic from field trees, enough leaf material was collected from individual PCR positive trees in the fields and stored at -80^oC in deep freeze and was used in subsequent studies. One negative control was kept.

3.3. DNA isolation from citrus tissue

Results (Table 2) revealed that the average DNA yield of three samples isolated by sodium sulphite method was 1216.16 ng/ µl. It was slightly higher in nucleic acid membrane method with average of 1429.16 ng/ µl. However, yield of DNA by kit method was only 159 ng/µl.

3.4. Determination of quality of DNA isolated by three methods

Results (Table 3) indicated that average value of DNA extracted by sodium sulphite method was 1.50 and by commercial kit was 1.54 while it was only 1.14 with DNA extractable by CTAB method.

The presence of greening disease in India was discovered by Fraser *et al.* (1966). Since then, the disease has been reported from various parts of the country (Ahlawat, 1997). In the absence of reliable diagnostic reagents and tools like electron microscope, the actual incidence and distribution of citrus greening bacterium (CGB) in India has not been investigated till 1991. During the period from 1991 to 1996, 51 and 339 trees from 98 orchards were tested which were collected from 8 different states of India and presence of CGB was confirmed by electron microscopy, immunofluorescence and DNA-DNA hybridization technique (Bove *et al.*, 1993, Ahlawat and Pant, 2003). Subsequently, the PCR technology was developed for detection of CGB (Harakava *et al.*, 2000; Hocquellet *et al.*, 2000; Tian *et al.*, 1996; Bove *et al.*, 1993; Jagouiet *et al.*, 1996). However this PCR technique was used for first time to detect greening disease in citrus in India by Ahlawat *et al.*, (2003). Serological-based diagnosis was not found as effective as it requires a panel of strain-specific monoclonal antibodies and due to sacrifice of animals in production of monoclonal antibodies. The nucleic acid hybridization technique required radioactive material and is not advisable if other efficient and reliable techniques like PCR are available. Therefore, during the present investigation, the information has been developed on suitability and reliability of PCR technique for detection of greening bacterium as a routine procedure of indexing. There was little information available of characterization of

Asian greening bacterium by amplification, cloning and sequencing although reports are three of African greening bacterium (Planet *et al.*, 1995)

Table 2

The quantity of DNA obtained from three infected sample by 3 method of DNA isolation

Samples	Sodium sulphite (ng/ μ l)	Commercial kit (ng/ μ l)	Nucleic acid (ng/ μ l)
1	1531.0	123.25	535.0
2.	2864.0	201.00	1056.0
3.	986.5	165.00	2113
4.	796.0	124.75	1053
5.	552.0	207.50	2212
6.	568.0	136.50	1606
Average	1216.16	159.60	1429.16

Table 3

Quality comparison of three methods (A_{260}/A_{280})

Samples	Sodium sulphite	Commercial kit	Nucleic acid membrane
1	1.06	1.05	1.14
2	1.53	2.86	1.02
3	1.13	1.05	1.01
4	2.54	1.44	0.99
5	1.76	1.60	1.55
6	1.01	1.27	1.15

During present study, trees of mandarin of the age of 3-10 years were periodically surveyed at the orchards of Marathwada region. A very high incidence of 65.00 per cent was observed. The greening incidence information in these investigation, was determined on the basis of typical symptoms of greening disease in field trees and the analysis of six candidate trees by polymerase chain reaction (PCR) using protocol. Six trees were identified based on PCR reaction for these studies. The leaves from these trees were collected and preserved at 80⁰C in deep freeze for the use in various experiments. The graft transmission was also obtained from this tree on Mosambi seedling after 3-6 months of grafting. However, only 4 plants out of six grafted, showed positive amplification in PCR. This may happen because the scions grafted on these two PCR negative plants may not be having the required quantity of pathogen required for graft transmission. It is known that the distribution of greening bacterium is erratic in the plant tissues (Varma *et al.*, 1993; Bove *et al.*, 1993) and this could be the reason of non transmission in plants by grafting. Since the studies were planned with a view to standardize and validate PCR diagnostics in field trees, the materials collected from six identified greening positive trees were used in these studies.

Three components are important for detection of pathogen in PCR. They are: a) Nucleic acid isolation and its quantity and quality b) Primer designing synthesis and its evaluation c) Evaluation of polymerase enzymes for PCR amplification.

During the present studies all the three steps were standardized. The DNA was isolated by three methods, sodium sulphate (Baranwal *et al.*, 2003). CTAB method (Murray and Thompson, 1980) and commercial kit obtained from QIAGEN Germany. The leaf material collected from 6 PCR positive trees was used in most of the experiments. The quantity of the DNA obtained from six samples was more CTAB method followed by Sodium sulphite method. Similar results were obtained by Baranwal *et al.* (2003) where commercial kit and CTAB method used while working with potato and a cherry virus and with citrus yellow mosaic virus. However, the yield of DNA by commercial kit (159.60 ng/ml) was much less as against sodium sulphate method (1216.16 ng/ml). The quality of DNA obtained by all the three methods was assessed by calculating A_{260}/A_{280}

ratio was observed that quality obtained by Sodium Sulphite method and commercial kit was almost at par (1.50 and 1.54, respectively). Although the best quality of DNA is known with the A_{260}/A_{280} of 1.8. However, this ratio has not been achieved with the any of the methods. But the satisfactory amplification was obtained even

of the DNA extracted by sodium sulphite method with the A_{260}/A_{280} ratio of 1.50. That is why this method was used in the experiment too. Since the quantity and quality of DNA isolated by sodium sulphite method was found to be better, this method was preferred over the methods of commercial kit and nucleic acid membrane method.

During the present studies, primers were not designed, but they were synthesized based on the published sequence data by Harakava *et al.*, (2000). Three parts of primers from conserved region of ribosomal β operon gene and 16 S rDNA were got synthesized by Sigma, Germany. These three pairs of primers were designated as A, B, C which provides amplicon size as 1160 bp, 703 bp and 450 bp, respectively. These three sets of primers were evaluated DNA. The comparative studies show that C sets of primer (450 bp) was superior than A and B as it could amplify the DNA upto 100 pg as against only 100 ng and 10 ng dilution of DNA with A and B sets of primers. The evaluation was also done using Taq and Klen Taq enzymes and Klen Taq as was expected gave better intensity of the bands obtained in 1 % agarose gel. Similar evaluation for primer has also been used by other researchers with different viruses (Singh and Nie, 2003; Harakava *et al.*, 2000).

During the present study, the DNA extraction method by sodium sulphite, Klen Taq enzyme and 450 bp set of primers were identified to give best amplification in PCR system of diagnosis. Using these protocols, samples from mandarin tree collected in March, 2010 were analyzed in PCR using combination of all the three standardized steps. It was found that the amplification of DNA isolated from out of twenty samples, 15 and 19 were PCR positive during the month of October and March, respectively. During the survey of greening diseases 65.00 per cent of incidence of disease was observed while testing trees for the presence of greening bacterium in PCR, 15 were positive during October and 19 in March indicating much higher incidence of disease with PCR testing. The results suggested that the PCR is reliable technique for detection of greening bacterium as against the diagnosis based on visible symptoms. It was therefore evident that protocols developed during present studies can detect greening bacterium in field trees irrespective of the season and the concentration of pathogen in the host. However, since more samples showed the presence of greening bacterium during March, the samples for detection of the bacterium should be taken in March to get maximum detection.

Detection of viruses and virus like pathogens as by PCR technique are gaining importance over the other technique as it is comparatively sensitive and reliable and can detect the pathogen even at a very low concentration upto 0.1 pg. The concentration of greening bacterium is very low in sieve tubes and distribution is also uneven (Varma *et al.*, 1993) and it is important that the samples are taken from different tree braches. The technology of PCR detection of greening bacterium developed during present investigation is new and validated for the first time in India. It will be very useful for field diagnosis of greening disease in planting material. The technology developed is cost effective and highly reliable for indexing bud wood certification programmes in citrus in India elsewhere. These studies will also be useful in plant quarantine for export and important of citrus germplasm.

References

- Ahlawat, Y.S., 1997. Viruses, greening, bacterium and viroids associated with citrus (*Citrus* sp.) decline in India. Indian J. Agril. Sci., 67 ,51-57.
- Ahlawat, Y.S., Pant, R.P., 2003. Major virus and virus-like diseases of citrus in India, their diagnosis and management. Annual Rev. Plant Pathol., 2 , 447-474.
- Ahlawat, Y.S., Baranwal, V.K., Mazumdar, S., 2003. First report of citrus greening disease and associated bacterium. Candidatus Liberibacter from Bhutan. Plant Disease, 87, 448.
- Baranwal, V.K., Majumdar, S., Ahlawat, Y.S., Singh, R.P., 2003. Sodium sulphite yields improved DNA of higher stability for PCR detection of citrus yellow mosaic virus from citrus leaves. J. of Virological methods, 112 , 153-156.
- Bove, J.M., Garnier, M., Ahlawat, Y.S., Chakraborty, N.K., Verma, A., 1993. Detection of Asian strains of greening BLO by DNA-DNA hybridization in Indian orchards trees and Malysian Diaphorina citri psyllids pp , 258-263 In Proc. 12th Conf. IOCV, IOCV Riverside, California.
- Fraser, L.R., Singh, D., Capoor, S.P., Nariani, T.K., 1966. Greening virus, the likely cause of citrus dieback in India. FAO Plant Prot. Bull., 14 , 127-130.
- Harakava, R., Morais, L.J., Ochasou, J., Manjunath, K.L., Febres, V.J., Lee, R.F., Niblett, G.L., 2000. Improved sensitivity in the detection and differentiation of citrus Huanglongbing bacteria from South Africa and the Phillippines. In. Proc., 14th Conf., IOCV. IOCV. Riverside, California. pp , 195-199

- Hocquellet, A., Bove, J.M., Garnier, M., 2000. Isolation of *Candidatus liberibacter* genes by RAPD and new PCR detection technique. In : Proc. 14th Conf. IOCV, IOCV Riverside, California. PP , 363-368
- Jagoueix, S., Bove , J.M., Grnier, M., 1996. PCR detection of the two candidates liberobacter species associated with greening diseases of citrus. Molecular and Cellular Probes, 10 , 43-50.
- Korsten, L.G., Sanders, M., Su, H.J., Garnier, M. Bove, J.M., Gotze, J.M., 1993. Detection of citrus greening infected citrus in South Africa using DNA probes and monoclonal antibodies. In Proc. 12th Conf. IOCV, IOCV, Riverside. pp , 224-232
- Planet, P.S., Jagoueix, B.J.M., Garnier, M., 1995. Detection and characterization of the African citrus greening Liberobacter by amplification, cloning and sequencing of rplKJL-ropBC operon. Curr. Microbiol. 30 , 137-141.
- Singh, R.P., Nie, X., 2003. Multiple virus and viriod detection and strain seperation via multiplex riverse transcription-polymerase chain reaction. Can J. Plant. Path., 25 , 127-134.
- Tian, Y., Ke, S., Ke, C., 1996. Polymerase chain reaction for detection and quantification of liberibacter asiaticus, the bacterium associated with Huanglongbing (Greening) of citrus in China. In Proc. 13th Conf., IOCV. IOCV Riverside, California. pp , 252-257
- Varma, A., Ahlawat, Y.S., Chakraborty, N.K., Garnier, M., Bove, J.M., 1993. Detection of greening BLO by electron Microscopy, DNA hybridization in citrus leaves with and without mottle from various regions in India. In Proc. 12th Conf., IOCV Riverside, California. pp , 280-285.