

Contents lists available at Sjournals



Journal homepage: www.Sjournals.com



Original article

Modeling and simulation to predict partial deposition of ammonia and specific growth rate of bacterial in homogeneous formation in Obio/Akpor, rivers state of Nigeria

S.N. Eluozo

Subaka Nigeria Limited Port Harcourt; Civil and Environmental Engineering Consultant Department of Research and Development, Rivers State of Nigeria.

*Corresponding author; Subaka Nigeria Limited Port Harcourt; Civil and Environmental Engineering Consultant Department of Research and Development, Rivers State of Nigeria.

ARTICLE INFO

Article history:

Received 23 January 2013

Accepted 18 February 2013

Available online 28 February 2013

Keywords:

Simulation

Partial deposition

Ammonia and homogenous formation

ABSTRACT

Specific growth rate of bacteria has been expressed through characteristics influence deposition at various stratum to aquiferous zone. The growth rates of bacteria are under the influence of the substrate utilization that contaminates groundwater aquifers in various formations. Derived mathematical equations generated theoretical values that expressed the rate of exponential phase of ammonia deposition and bacteria growth rate in homogenous formation. Slight fluctuations were experienced in between the aquiferous zones under the influence of variation from degree of porosity and permeability of the stratum. Experimental values were compared with the theoretical values; both parameters produced a best fit. These expressed the application of the derived model that determines partial deposition of ammonia and bacteria growth rate in homogenous formation. The study is imperative because the model expressed partial depositions of ammonia that determine the growth rate of bacteria at various formations to groundwater aquifers in the study area.

© 2013 Sjournals. All rights reserved.

1. Introduction

Ammonia is used in fertilizer and animal feed production and in the manufacture of fibers, plastics, explosives, paper, and rubber. It is used as a coolant, in metal processing, and as a starting product for many nitrogen-containing compounds (3). Ammonia and ammonium salts are used in cleansing agents and as food additives (1,4), and ammonium chloride is used as a diuretic [Source: *Hazardous Substances Data Bank: Ammonium chloride*. Bethesda, MD, National Library of Medicine, 1990, WHO 1996]

Microelements are nutritional components that occur and function in many locations and on many levels of organism. They are present in the organism at very low concentrations and are an indispensable component in numerous enzymatic, catalytic, regulating and activating processes, usually as activators and co-factors. Microelements enter the organism through the feed; in developing foetuses they enter through the placenta. Concentration of microelements in blood of supplemented mothers correlates positively with concentrations in calf blood. An important source of microelements for a newborn calf is the colostrums (Abdelrahman and Kincaid 1993; Lacetera 1996; Underwood and Suttle 1999; Pavlata et al. 2003). Microelements also affect the quality and composition of the colostrums and milk and affect the health of the udder. For example, the colostrums and milk of selenium-supplemented cows has a higher concentration of selenium and contains a higher concentration of immunoglobulin's; cows supplied with selenium have lower incidence of mastitis (Hogan et al. 1993; Knowles et al. 1999; Pavlata et al. 2004a); supplementation of zinc to dairy cows decreases the number of somatic cells in milk (Pechová et al. 2006), and copper affects the ability of neutrophils to kill phagocytosed bacteria and decreases susceptibility of the udder to infection (Scaletti et al. 2003). Microelement deficiencies in dairy and beef cattle in the Czech Republic are frequent (Pavlata et al. 2005a; Slavík et al. 2006; Podhorský et al. 2007). Ruminant nutrition routinely uses several methods and forms to supplement microelements. A slightly problematic period for ensuring adequate supplementation of microelements to dairy cows is the dry period, when only a small amount of seeds is usually fed and microelements are thus difficult to add to the feed. Application forms include supplementation by adding minerals to the feed, mineral licks, and/or single/repeated injection of individual microelements, or combinations of the above. Other methods of supplementation include boluses, projectiles or pills (depending on the shape of administration form), containing a precisely defined amount of microelements combined with a carrier or auxiliary substance. Another method of supplementation of microelements consists in long-acting injections (Lee et al. 1999; Pavlata 2004a; Kinal et al. 2004; Pechová et al. 2006; Mulligan et al. 2006; Chládek and Zapletal 2007).

2. Materials and Methods

Analytical model were developed, applying mathematical tools, the derived model were applied produced theoretical values that were compared with experimental laboratory analysis. The experimental procedure is column experiment, the soil samples were collected at intervals of three metres each (3m). Ammonia solute was introduced at the top of the column and effluents from the lower end of the column were collected and analyzed for Ammonia that generated results from its analysis

2.1. Governing Equation

$$\frac{\mu_o}{\mu_n} \frac{\partial K_o}{\partial t} = \frac{\partial K_o}{\partial x} K_v + K_c \quad \dots\dots\dots (1)$$

Nomenclature

- θ = Void Ratio
- K = Permeability
- V = Velocity
- T = Time
- X = Distance

h_{AO} = Concentration deposition phosphorus
 Kc = Inhibitors of substrate

$$K_o = XT$$

$$\frac{\partial K_o}{\partial t} = XT^1 \dots\dots\dots (2)$$

$$\frac{\partial K_o}{\partial X} = TX^1 \dots\dots\dots (3)$$

$$\frac{\mu_o}{\mu_n} \frac{T^1 X}{TX} = \frac{TX^1}{TX} K_v + K_c = -\lambda^2 \dots\dots\dots (4)$$

$$\frac{\mu_o}{\mu_n} \frac{T^1 X}{TX} = \frac{X^1}{X} K_v + K_c = -\lambda^2 \dots\dots\dots (5)$$

$$\frac{\mu_o}{\mu_n} \frac{T^1}{T} = -\lambda^2 \dots\dots\dots (6)$$

$$\frac{X^1}{X} K_v + K_c = -\lambda^2 \dots\dots\dots (7)$$

From (7) $T^1 + \frac{\lambda^2 T}{\frac{\mu_o}{\mu_n}} = 0$

$$X = A \cos \frac{\lambda}{\sqrt{\frac{\mu_o}{\mu_n}}} t + B \sin \frac{\lambda}{\sqrt{\frac{\mu_o}{\mu_n}}} t \dots\dots\dots (8)$$

From (6) $\frac{Z^1}{Z} K_v + K_c - \lambda^2 \dots\dots\dots (9)$

$$\frac{Z^1}{Z} = \frac{-\lambda^2}{K_v K_c} \dots\dots\dots (10)$$

By direct integration

$$\ln T = \frac{\lambda^2}{K_v K_c} Z \dots\dots\dots (11)$$

$$Z = D \ell^{\frac{-\lambda^2}{K_v K_c} Z} \dots\dots\dots (12)$$

Combining (8) and (9) yields

$$K_o(Z,t) = \left[A \cos \frac{\lambda}{\sqrt{\mu_o}} t + B \sin \frac{\lambda}{\sqrt{KvKc}} Z \right] D \ell^{\frac{-\lambda^2}{KvKc} Z} \dots\dots\dots (13)$$

At $t = 0$ $K_o(o) = K_o$

$$K_o = A \cos \frac{\lambda}{\sqrt{\mu_o}} t + B \sin \frac{\lambda}{\sqrt{KvKc}} Z$$

$$\frac{\partial K_o}{\partial t} \Big|_{t=0} = 0 \dots\dots\dots (14)$$

From (13) $\frac{\partial K_o}{\partial t} = \left[-A \frac{\lambda}{\sqrt{\mu_o}} \sin \frac{\lambda}{\sqrt{\mu_o}} t + B \frac{\lambda}{\sqrt{\mu_o}} \cos \frac{\lambda}{\sqrt{\mu_o}} t \right] D \ell^{\frac{-\lambda^2}{KvKc} Z} \dots\dots\dots (15)$

At $t = 0$

$$0 = B \frac{\lambda}{\sqrt{KvKc}} D \ell^{\frac{-\lambda^2}{KvKc} Z} \Rightarrow B = 0 \quad D \neq 0 \dots\dots\dots (16)$$

$$K_o = \left[A \cos \frac{\lambda}{\sqrt{\mu_o}} \right] D \ell^{\frac{-\lambda^2}{KvKc} Z} \dots\dots\dots (17)$$

$$K_o = AD \cos \frac{\lambda}{\sqrt{\mu_o}} t \ell^{\frac{-\lambda^2}{KvKc} Z} \dots\dots\dots (18)$$

$$\frac{\partial K_o}{\partial t} = AD \frac{-\lambda}{\sqrt{\mu_o}} \sin \frac{\lambda}{\sqrt{\mu_o}} t \ell^{\frac{-\lambda^2}{KvKc} Z} \dots\dots\dots (19)$$

At $t = \frac{\partial K_o}{\partial t} = 0$

$$0 = \frac{AD\lambda}{\sqrt{\frac{\mu_o}{\mu_n}}} \text{Sin} \frac{\lambda d}{\sqrt{\frac{\mu_o}{\mu_n}}} = n\pi = \frac{\lambda d}{\sqrt{\frac{\mu_o}{\mu_n}}}, n=0,1,2 \dots\dots\dots (20)$$

$$\Rightarrow \lambda = n\pi \frac{\sqrt{\frac{\mu_o}{\mu_n}}}{d} \dots\dots\dots (21)$$

So that we have

$$K_o (Z,t) = AD \text{Cos} n\pi \frac{\sqrt{\frac{\mu_o}{\mu_n}}}{d} t \ell \frac{-n^2 \pi^2 \frac{\mu_o}{\mu_n}}{d^2 K_v K_c} Z \dots\dots\dots (22)$$

$$AD \text{Cos} \frac{n\pi}{d} t \ell \frac{-n^2 \pi^2 \frac{\mu_o}{\mu_n}}{d^2 K_v K_c} Z \dots\dots\dots (23)$$

Hence $AD = K_o$

$$K_o = (Z,t) = K_o \ell \frac{-n^2 \pi^2 \frac{\mu_o}{\mu_n}}{d^2 K_v K_c} Z \text{Cos} \frac{n\pi}{d} t$$

\dots\dots\dots (24)

3. Results and discussion

Modeling and simulation to predict partial deposition of ammonia and specific growth rate of bacterial in homogeneous formation in are presented in tables and figures bellow.

Table 1
Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.

Depth M	Theoretical values	Experimental values
3	1.43E-06	1.48E-06
6	2.86E-06	2.77E-06
9	4.29E-06	4.44E-06
12	5.71E-06	5.67E-06
15	6.35E-06	6.44E-06
18	7.62E-06	7.88E-06
21	8.89E-06	8.56E-06
24	1.02E-05	1.11E-05
27	1.14E-05	1.18E-05
30	1.27E-05	1.32E-05

Table 2

Comparison of theoretical and experimental values of partial deposition of ammonia at various times.

Time per Day	Theoretical values	Experimental values
10	1.43E-06	1.48E-06
20	2.86E-06	2.77E-06
30	4.29E-06	4.44E-06
40	5.71E-06	5.67E-06
50	6.35E-06	6.44E-06
60	7.62E-06	7.88E-06
70	8.89E-06	8.56E-06
80	1.02E-05	1.11E-05
90	1.14E-05	1.18E-05
100	1.27E-05	1.32E-05

Table 3

Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.

Depth M	Theoretical values	Experimental values
3	2.84E-06	2.77E-06
6	1.70E-05	1.66E-05
9	2.55E-05	2.44E-05
12	3.40E-05	3.45E-05
15	4.26E-05	4.45E-05
18	5.11E-05	5.55E-05
21	5.96E-05	6.11E-05
24	6.81E-05	6.55E-05
27	7.67E-05	7.44E-05
30	8.52E-05	8.44E-05

Table 4

Comparison of theoretical and experimental values of partial deposition of ammonia at various times.

Time per Day	Theoretical values	Experimental values
10	2.84E-06	2.77E-06
20	1.70E-05	1.66E-05
30	2.55E-05	2.44E-05
40	3.40E-05	3.45E-05
50	4.26E-05	4.45E-05
60	5.11E-05	5.55E-05
70	5.96E-05	6.11E-05
80	6.81E-05	6.55E-05
90	7.67E-05	7.44E-05
100	8.52E-05	8.44E-05

Table 5

Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.

Depth M	Theoretical values	Experimental values
3	5.96E-08	5.88E-07
6	2.38E-07	2.45E-07
9	5.11E-07	5.33E-07
12	9.53E-07	9.44E-07
15	1.59E-06	1.64E-06
18	2.14E-06	2.23E-06
21	2.90E-06	2.88E-06
24	3.81E-06	3.77E-06
27	4.80E-06	4.77E-06
30	5.96E-06	5.88E-06

Table 6

Comparison of theoretical and experimental values of partial deposition of ammonia at various times.

Time per Day	Theoretical values	Experimental values
10	5.96E-08	5.88E-07
20	2.38E-07	2.45E-07
30	5.11E-07	5.33E-07
40	9.53E-07	9.44E-07
50	1.59E-06	1.64E-06
60	2.14E-06	2.23E-06
70	2.90E-06	2.88E-06
80	3.81E-06	3.77E-06
90	4.80E-06	4.77E-06
100	5.96E-06	5.88E-06




Fig. 1. Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.




Fig. 2. Comparison of theoretical and experimental values of partial deposition of ammonia at various times.




Fig. 3. Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.




Fig. 4. Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.




Fig. 5. Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.




Fig. 6. Comparison of theoretical and experimental values of partial deposition of ammonia at various depths.

From figure one to six the deposition of ammonia are found to be in exponential phase as presented in the figures, the growth rate of ammonia shows that the deposition of other influences generated the fast deposition of ammonia as presented in the figures. Slight fluctuations were experienced in between the fine and coarse sand at forty to sixty days within twelve and eighteen metres on both parameters of theoretical and experimental values. Porosity and permeability expressed their influence based on high degree of depositions between the silt strata formation to coarse formation from twelve metres to thirty metres. This influence generated the migration with high concentration in some figures, but at the aquiferous zone, it expresses low concentration which may be insignificant on groundwater quality, unless there is regeneration as expressed in the figures presented. The deposition of ammonia implies that areas where there is high concentration will definitely generate increase of microbial population that will contaminate groundwater aquifers in the study location. The expression of ammonia deposition from the developed model shows that the rates of concentration coefficient of inhibition were insignificant as inhibitors deposition were found to be very low at various formations. Results at aquiferous zone can be compared to World Health Organization to determine its quality for human consumption. The study is imperative because the model can be applied to monitor partial depositions of ammonia within the influence of coefficient inhibition as it reflects fast migration of microbes to groundwater aquifer. Practicing engineers, scientists and public health officers will use this concept to monitor the growth rate of microbial contaminant in groundwater aquifers.

4. Conclusion

Partial deposition of ammonia and specific growth rate of bacterial in homogenous formation has been expressed. Expressed equations were derived and it generated a model equation for partial deposition of ammonia and specific growth rate of bacterial in homogenous formation. This expression generated theoretical values that show the exponential phase of ammonia with respect to growth rate of bacterial at various formations. Formation characteristics were found to influence the deposition of ammonia at various formations down to groundwater aquifers. Experimental results were compared with theoretical values and it expressed a best fit in all the figures presented, this implies that the model can be applied to monitor the partial deposition of ammonia and growth rate of bacterial on homogenous formation. Geologic history expressed alluvium deposition found to be

homogeneous formation generated a homogenous soil stratum that expressed its influence on the partial deposition of ammonia and specific growth rate in the study location. The study is imperative because it has shown how homogeneous formations play a major role on partial deposition of ammonia and specific growth rate of bacteria in homogeneous stratum.

References

- Chládek, G., Zapletal, D., 2007. A free-choice intake of mineral blocks in beef cows during the grazing season and in winter. *Livest Sci* 106, 41-46
- World Health Organization, 1996. Originally published in Guidelines for drinking-water quality, 2nd ed. Vol. 2. *Health criteria and other supporting information*.
- Abdelrahman, M.M., Kincaid, R.L., 1993. Deposition of copper, manganese, zinc, and selenium in bovine fetal tissue at different stages of gestation. *J Dairy Sci* 76, 3588-3593
- Lacetera, N., Bernabucci, U., Ronchi, B., Nardone, A., 1996. Effects of selenium and vitamin E administration during a late stage of pregnancy on colostrum and milk production in dairy cows, and on passive immunity and growth of their offspring. *Am J Vet Res* 57, 1776-1780
- Pavlata, L., Pechová, A., Dvořák, R., 2004b. Microelements in colostrum and blood of cows and their calves during colostrum nutrition. *Acta Vet Brno* 73, 421-429
- Pavlata, L., Pechová, A., Illek, J., 2000. Direct and indirect assessment of selenium status in cattle - a comparison. *Acta Vet Brno* 69, 281-287
- Pavlata, L., Podhorský, A., Pechová, A., Chomát, P., 2005a. Differences in the occurrence of selenium, copper and zinc deficiencies in dairy cows, calves, heifers and bulls. *Vet Med-Czech* 50, 390-400
- Pavlata, L., Prášek, J., Podhorský, A., Pechová, A., Haloun, T., 2003. Selenium metabolism in cattle: maternal transfer of selenium to newborn calves at different selenium concentrations in dams. *Acta Vet Brno* 72, 639-646
- Pavlata, L., Prášek, J., Filípek, J., Pechová, A., 2004a. Influence of parenteral administration of selenium and vitamin E during pregnancy on selected metabolic parameters and colostrum quality in dairy cows at parturition. *Vet Med-Czech* 49, 149-155
- Pechová, A., Pavlata, L., Hofírek, B., Dvořák, R., 2004. Metabolic profile test. In: Hofírek B, Pechová A, Doležel R, Pavlata L, Dvořák R, Fleischer P et al. (Eds): *Herd health management in dairy cattle (in Czech)*. VFU Brno, pp. 31-49
- Podhorský, A., Pechová, A., Dvořák, R., Pavlata, L., 2007. Metabolic disorders in dairy calves in postpartum period. *Acta Vet Brno* 76, 45-53.
- Scaletti, R.W., Trammell, D.S., Smith, B.A., Harmon, R.J., 2003. Role of dietary copper in enhancing resistance to *Escherichia coli mastitis*. *J Dairy Sci* 86, 1240-1249.
- Lee, J., Masters, D.G., White, C.L., Grace, N.D., Judson, G.J., 1999. Current issues in trace element nutrition of grazing livestock in Australia and New Zealand. *Aust J Agric Res* 50, 1341-1364
- Slavík, P., Illek, J., Zelený, T., 2006. Zinc and copper status of beef cattle in the Šumava region, Czech Republic. *Acta Vet Brno* 75, 485-488.
- Mulligan, F., O'Grady, L., Rice, D., Doherty, M., 2006. A herd health approach to dairy cow nutrition and production diseases of the transition cow. *Anim Reprod Sci* 96, 331-353.
- Kinal, S., Rzasna, A., Korniewicz, A., 2004. Mineral bioplex supplementation of diets for cows affects colostrums quality and immunoglobulins in calf blood serum. *J Anim Feed Sci* 13 (Suppl.), 79-82.
- Krys, Š., Lokajová, E., Podhorský, A., Pavlata, L., 2009. Microelement Supplementation in Dairy Cows by Mineral Lick *acta vet. brno* 78, 29-36; doi:10.2754/avb200978010029.