

Provided for non-commercial research and education use.

Not for reproduction, distribution or commercial use.



This article was published in an Sjournals journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the authors institution, sharing with colleagues and providing to institution administration.

Other uses, including reproduction and distribution, or selling or licensing copied, or posting to personal, institutional or third party websites are prohibited.

In most cases, authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Sjournals's archiving and manuscript policies encouraged to visit:

<http://www.sjournals.com>

© 2018 Sjournals Publishing Company

Contents lists available at Sjournals
Scientific Journal of Veterinary Advances

Journal homepage: www.Sjournals.com



Original article

Preliminary study on small ruminant GIT helminthiasis in select arid and semi-arid pastoral and agro-pastoral areas of Afar region, Ethiopia

Hailegebrael Bedada^{*}, Fikru Gizaw, Wossene Negash

College of Veterinary Medicine, Samara University, P.O. Box 132, Semera, Afar, Ethiopia.

*Corresponding author: hailegebraelbedada@su.edu.et

ARTICLE INFO

Article history,

Received 09 December 2017

Accepted 10 January 2018

Available online 17 January 2018

iThenticate screening 11 December 2017

English editing 09 January 2018

Quality control 16 January 2018

Keywords,

Helminths

Small ruminants

Prevalence

Afar region

Ethiopia

ABSTRACT

A cross-sectional study was conducted to determine the prevalence and identification of GIT parasites of sheep and goats in two selected districts of Afar from December 2014 to February 2016. Totally 590 faecal samples were collected from small ruminants (332 goat and 258 sheep) managed in pastoral and agro-pastoral production. Out of the total examined small ruminant 87.8% (518) were found to harbor one or more genera of helminth parasites. The result of the study revealed that 92.2% (306) of the goats and 82.2% (212) of the sheep were found positive for GIT parasites. Helminth parasites identified in small ruminant of the study area were Strongyles, Fasciola, Strongyloides, Paramphistomum, Trichuris, Ascaris and Monezia. The risk of infection with GIT helminth parasites in goats were 4.009 times higher than sheep (OR=4.009, p=0.011). Age and sex related difference was not observed in the prevalence of helminth parasites in sheep and goats. Significantly (OR=0.119, p=0.000) higher prevalence of overall helminth parasites in poor body condition sheep and goats than good body condition was observed. Likewise, significant variation in overall parasite prevalence was observed between the study districts (OR=0.169, p=0.000). In this study, species of the animals, origin, and body condition score are important risk factors associated with gastrointestinal parasites in the study area. In the study area nutrition is generally poor, low productivity in small ruminants is likely to be aggravated by a high prevalence of polyparasitism.

© 2018 Sjournals. All rights reserved.

1. Introduction

Sheep and goat have an enormous potential to affect the socioeconomic development of the majority of African rural communities. Increasing small ruminant production can boost farm income by generating cash income that can be used to purchase inputs for other production activities hence improves the quality of life of the people of the Sub-Saharan Africa (UNECA, 2012). Small ruminants account the most important part of the Ethiopian livestock system, about 24.2 million sheep and 22.6 million of goats reported in the country. In Ethiopia small ruminants are exist in all agro climatic zones. The highland area comprises 75% of the sheep and 27% of the goat population, while the lowland pastoral and agro pastoral area have 25% of the sheep and 73% of the goats' population (CSA, 2012). In pastoralist area, goats and sheep are mainly utilized for milk and meat production and generate income to the owner. Nearly 87% of the Afar regional state population are rural mainly dependent on pastoral and agro-pastoral livelihood systems. In the arid areas of Afar, which are none or limited agricultural potential livestock production is the main and only possible activity of the pastoralists. Among the livestock sectors on which the pastoralists of the region make their livelihood, small ruminant are the main cash income and play an important role. In spite of the large population and potential use of small ruminants in pastoral area, the production system is affected by feed shortage, poor genetic makeup of the animals, and wide spread occurrence of livestock diseases such as endoparasites which have great economic significance to the communities and the country as a whole (FAO, 2005).

Health disorders in all classes of small ruminants represent a factor that greatly affects the economics of sheep and goat production. The most serious problem confronting sheep and goats production worldwide is infection with GIT parasites. Parasitic infection ranks high among the factors that interfere the productivity of small ruminants although its effect is often underestimated. Correspondingly the current levels of contributions of the small ruminants in Ethiopia, either the macro or micro level is lower than the expected potential. Among major constraints hindering the productivity of sheep and goat in the country diseases are the principal one; of which endoparasites play a wide range of health problems that tackle the productivity of small ruminants (Chanie and Begashaw, 2012). In Ethiopia, parasitological investigations of small ruminants in the highland parts of the country have demonstrated that GIT parasites are the most common problem affecting production and productivity of the area sheep and goats. However, study on the GIT helminth parasites of small ruminants in the Afar region is not yet conducted. Likewise, in the study area different animal species and different age group were kept together on the same pasture and all species of animals share common water point. Hence, such type of husbandry system endorsed transmission and maintenance of parasites. Therefore, the objective of this study was investigating GIT helminth parasites infecting small ruminants of the study area.

2. Materials and methods

2.1. Study area and population

The study was conducted in two districts (Asayita and Dubti) of Afar pastoral and agro-pastoral region. The Afar National Regional State is located in the Great Rift Valley, comprising rangeland in northeast Ethiopia with an estimated area of 95,958 Km². The Afar region is located in Northeast of Ethiopia between 39°34' and 42°28' E longitude and 8°49' and 14°30' N latitude. In the Afar region, there are about 4,268,000 goats and 2,464,000 sheep, 2,336,488 heads of cattle, 852,016 camels and 187,287 equines which are managed under pastoral and agro-pastoral production system. The annual temperature and rainfall in the region is 30-50°C and 200-600mm, respectively. The altitude of the region ranges from 116 meter below sea level to 1600 meters above sea level (CSA, 2012).

Dubti district contained about 49,234 goats, 24,363 sheep, 39,412 heads of cattle, 7241 camels and 2295 equines which are managed under pastoral and agro-pastoral production system (CSA, 2012). The production system in the area predominantly is pastoralism and agro-pastoralism. The major feed resource in the district is natural grazing and supplements from maize, sorghum, cotton and sesame. Asayita district is located in between 110°34' N and 410°26' E. The annual temperature and rainfall in the district is 28-41.7°C and 144mm annual precipitation, respectively. Asayita district located at an elevation of 340 meter above sea level (ANRS, 2010). In Asayita district there are about 97,013 goats, 17,198 sheep, 81,767 heads of cattle, 6108 camels and 3303 equines

which are managed under pastoral and agro-pastoral production system (CSA, 2012). The major feed resource in the district is natural grazing and supplements from maize, sorghum, cotton and sesame.

2.2. Study design and sampling methods

A cross-sectional study was employed from December 2014 to February 2016 to address the objective of the study. The sampling method was supposed to be a multi-stage cluster sampling approach. However, due to the absence of between cluster variance and sampling frame in the study districts and pastoral community as a whole and, during sample collection due to migration of animals, unwillingness of pastoralists to include their animals in this study, conflict among pastoralists therefore, the flocks of small ruminants were sampled purposively. However, proportional allocation was used to distribute the individual sampled small ruminants evenly among the flock. Consequently, 5 to 15 individual small ruminants from each small ruminants flock were sampled. Zone one was purposively selected based on the small ruminant population and topography. The primary stage was sampling of districts from the selected zone. Selection of Kebeles/PAs, flock and individual small ruminant within the flock were the 2nd, 3rd and 4th stages, respectively. Accordingly, two districts and four Kebeles/PAs from each district, proportional flocks from each Kebeles and 5-15 small ruminant population from each flock was sampled. Each Kebeles/PAs and individual animals from the flock sampled randomly. Therefore, total of 590 small ruminants was included in this study.

2.3. Sample collection

Fresh fecal samples approximately 10g were collected directly from the rectum of small ruminant and sample placed in sampling bottles (without preservatives) and labeled. Samples were transported to laboratory with ice box contain filled ice bag hence, there were no significant changes in the egg morphology. Following transportation of fecal sample laboratory analysis was carried out at the same day and the remaining samples were kept under 4°C and examined up on the next days. In the laboratory, fecal samples were examined for detection of helminth eggs using standard procedures of flotation and sedimentation methods; eggs will be identified based on their color, shape and contents (Soulsby, 1982; Hansen and Perry, 1994; Gareth, 2009). The collected fecal samples were processed and examined under the 10x and 40x magnification.

3. Results and discussion

Out of 590 small ruminants (332 goat and 258 sheep) examined 518 (87.8%) were infected with single or multiple parasites. From the totally examined small ruminant 92.2% goat and 82.2% sheep were infected with single or multiple parasites. The major helminth parasites identified from the small ruminants of study area were 49.2% *Strongyles*, 32.5% *Fasciola*, 28.5% *Strongyloides*, 25.4% *Trichuris*, 20.3% *Paramphistomum*, 8.8% *Ascaris* and 8.8% *Monezia* (Table 1 and Fig. 1). From 87.8% of small ruminants infected, 58.3% and 29.5% were infected by multiple parasites and single parasite respectively.




Fig. 1. GIT parasites identified in the study area.

Table 1

GIT parasites identified in the study area.

Parasites	Goat (n=332)	Sheep (n=258)	Total (n=590)
Trichuris	23.5%(78)	27.91%(72)	25.4%(150)
Ascaris	11.45%(38)	5.4%(14)	8.8%(52)
Strongyles	56.02%(186)	40.3%(104)	49.2%(290)
Strongyloides	31.33%(104)	24.8%(64)	28.5%(168)
Fasciola	37.95%(126)	25.58%(66)	32.5%(192)
Paramphistomum	24.1%(80)	15.5%(40)	20.3%(120)
Monezia	9.04%(30)	8.53%(22)	8.8%(52)
Overall	92.2%(306)	82.2%(212)	87.8%(518)

Statistically result revealed significant variation was recorded in the overall prevalence of parasitism between species (OR=4.009, $p=0.011$), body condition score (OR=0.119, $p=0.000$) and origin (OR=0.169, $p=0.000$) of the study animals (Table 2).

Table 2

Statistical result of overall helminth parasites infection of the study area.

Risk factors	SE	P-value	OR	95% CI for OR	
				Lower	Upper
Species	0.549	0.011	4.009	1.368	11.751
Age	0.457	0.109	2.081	0.850	5.097
Sex	0.519	0.571	0.745	0.270	2.062
BCS	0.486	0.000	0.119	0.046	0.308
Origin	0.495	0.000	0.169	0.064	0.447

3.1. Prevalence by the origin of small ruminants

The overall prevalence of small ruminant parasite was significantly higher in Asayita district (93.13%) than in Dubti district (81.5%) (OR=0.169, $p=0.000$) (Table 1). However, the result showed high parasitic infection in both districts.

3.2. Species based prevalence

The overall prevalence of GIT parasites in goat and sheep of the study area was 92.2% and 82.2% respectively. Statistically significant variation in the overall prevalence of parasites was recorded between goat and sheep of the study area (OR=4.009, $p=0.011$) higher in goat than sheep (Table 1).

3.3. Prevalence of parasite by sex

Prevalence of GIT parasites in female and male animal of the study area was 88.13% and 87.16% respectively. Significant difference in the prevalence of parasites was not recorded between male and female small ruminants of the study area ($p>0.05$) (Table 1).

3.4. Prevalence of parasites by age

Statistically significant difference was never recorded ($p>0.05$) in the overall prevalence of parasite infestation between young and adult small ruminants of the study (Table 1). Prevalence of GIT parasites in young and adult small ruminants of the study area was 83.78% and 90.2% respectively (Table 1).

3.5. Prevalence of parasites by body condition

The overall prevalence of parasites in good and poor body condition small ruminant of the study area were 75.22% and 95.6% respectively. The logistic regression results showed statistically significance difference in overall prevalence of parasites (OR=0.119, $p=0.000$) between small ruminants with poor and good body condition in study area (Table 1).

Results presented in this study revealed 87.8% an overall prevalence of small ruminant GIT helminthiasis in the study area. The current finding is in line with previous reports from different parts of Ethiopia and other tropical countries (Gebreyesus, 1986; Bikila et al., 2013). However this finding is much higher than the findings in different part of the country (Tesfaheywet, 2012; Temesgen, 2008). The high prevalence observed in this study most probably attributable to several important factors including management problems (keeping different species of animals and different age group together), conducive environment, malnutrition and negligible knowledge of pastoralists and scarce veterinary services in the study region. In addition due to no policy which limit free movement of animals from one place to the other in the country as a whole hence, clinically affected and carrier animals can serve as the source of infection. The effects of climate change and other factor such as changes in animal management and husbandry systems, usage of antiparasitics and increase in animal trafficking may also contribute to the changes in the prevalence, or emergence of parasitic infection in certain localities (Radiostits et al., 2007).

The overall prevalence of GIT parasite was significantly higher in goat than sheep of the study area ($p=0.011$). Goats of the study area were 4.009 times at risk for GIT infestation than sheep. The current finding is in agreement with works of different researchers (Fikru et al., 2006; Dagnachew et al., 2011). But, this finding is in contrary with other findings revealed GIT parasite infection is higher in sheep than goat (Bikila et al., 2013; Tesfaheywet, 2012; Teklye, 1991; Waruiru et al., 2005). Nonetheless, in the present finding even though sheep and goats differ in their feeding habits, small ruminants are kept together on common grazing land which may expose goats to acquire more susceptibility for the same species of GIT parasites. In consequence, the condition could be due to less or slow development of immunity in goats to GIT parasites compared with the situation in sheep. Goats do not build up an effective immune response against helminth infections and so remain susceptible to disease throughout their lives. The risk is enhanced if they are forced to graze rather than browse (Radiostits et al., 2007; Urquhart et al., 1996). Sheep faced prolonged challenge over generations and had developed good resistance (Urquhart et al., 1996). So that, in the study area recurrent drought occurred repeatedly, so goat of the study area forced to graze as the bush become dry during drought time. In general Afar pastoralists own different species of domestic animals, and these animals share common watering points and grazing pasture. The watering points of small ruminants are commonly shared with cattle and camel creating a close interspecies interaction among these domestic animals, and this might increases the risk of transmission of GIT parasites which infect multiple species.

Prevalence of helminth parasite is higher in adult animal than young one. However, age wise observation revealed no statistically significant difference in infestation of helminth parasites. This finding agrees with reports from Gambia and semi-arid part of Kenya indicated that GIT helminthes affect both ages insignificantly (Waruiru et al., 2005; Fritsch et al., 1993; Soulsby, 1982). The present finding disagrees with most literatures and studies (Fikru et al., 2006; Gamble and Zajak, 1992; Colditz et al., 1996) that young sheep and goats are more susceptible to parasite infection than adult small ruminants. The researchers justified the result that it could be because adult animals may acquire immunity to the parasite through frequent challenge and expel the ingested parasite before they establish infection and young animals are susceptible due to immunological immaturity and immunological unresponsiveness. However, in this study, we ascribe the absence of significant difference in parasites infection between young and adult animals might be due to the small number of young animals kept by the pastoralists as they interested to sale young animals since the community used small ruminant milk as a source of food for the household and if they keep young they may not fulfill their necessitate of using milk. Therefore, young small ruminant which are kept as replacement stock may enforced to graze as early age because of limited access to milk.

In the current study significant higher prevalence of gastrointestinal parasites were recorded in poor body condition small ruminant than good body condition animals, this finding agrees with (Temesgen, 2008; Keyyu et al., 2003; Ahmed, 1988; Sissay, 2007). In addition, Radiostits et al. (2007) and Odoi et al. (2007) showed that animals with poor condition are highly susceptible to infection and may be clinically affected by worm burdens too small to harm an otherwise well-fed healthy animal. Moreover, Kumba et al. (2003) observed that a well-fed animal was not in trouble with worms, and usually a poor diet resulted in more helminth infections. Furthermore, helminths also led to a loss of appetite and poor utilization of food, which results in a loss of body weight.

Fecal examination in small ruminants have shown the presence of *Strongyles*, *Trichuris*, *Strongyloides*, *Ascaris*, *Fasciola*, *Paramphistomum*, *Monezia* and mixed infections. The helminth parasites recorded in the study area have also been reported previously in other areas of Ethiopia (Bersisa et al., 2011; Fikru et al., 2006; Abebe et al., 2010; Hailelul, 2002; Tesfaye, 1998) and elsewhere in the world (Odoi et al., 2007; Kumba et al., 2003; Githigia

et al., 2001; Ntonifor et al., 2013). The current study showed that nematodes are the most common helminth parasites of both sheep and goats of the study area. The result also revealed that strongyle nematodes were identified as the most predominant helminths in the small ruminants of the study area. This finding is in agreement with different findings (Bersisa et al., 2011; Bikila et al., 2013; Tesfaheywet, 2012; Bashir et al., 2012; Sissay, 2007). The high prevalence of strongyles may be due to the suitability of the climatic condition of the study area for survival and transmission of the parasites. In addition, the poor management practices, including the poor hygienic practices employed by the pastoralists and keeping different age group and different species of animals together may be the other contributing factor as the epidemiology of nematodosis is determined by environmental factors (Radiostits et al., 2007; Urquhart et al., 1996; Soulsby, 1982).

4. Conclusion

The overall prevalence of GIT parasites in the study area indicates it is important health problem due to high prevalence and occurrence of polyparasitism. The prevalence was significantly influenced by species, origin of the animal and body condition score of the small ruminants. In the study area pastoralists own different species of domestic animals, and these animals share common watering points and grazing sites which creating a close interspecies interaction among these domestic animals, and this might increase the risk of transmission of GIT parasites which infect multiple species. In addition, shortage of feed source in the study area magnified the occurrence and prevalence of GIT parasites. Pastoralists of the study area are at risk of economic losses from decreased productivity of their animals as a majority of them harbored polyparasitism. Thus, GIT parasites should considered among those diseases responsible for health and productivity problems in small ruminant. More detailed studies on helminths should be conducted to pinpoint appropriate time for strategic deworming. In addition year rounded investigation is needed to know the species composition, survival strategy and ecology of the economically important GIT helminth parasites of sheep and goat in the study area.

Acknowledgements

The authors are grateful to Samara University, Semera Regional Veterinary Laboratory and individuals involved for the success of this work.

References

- Abebe, R., Gebreyohannes, M., Mekuria, S., Abunna, F., Regassa, A., 2010. Gastrointestinal nematode infection in small ruminant under traditional husbandry system during dry season in Southern Ethiopia. *Trop. Anim. Health*, 42, 1111-7.
- Afar National Regional State (ANRS), 2010. Programme of plan on adaptation to climate change, report, Semera. <http://ebookbrowse.com/afar-national-regional-stateclimate-change-adaptation-program-pdf-d127720299>
- Ahmed, N., 1988. Prevalence of GIT helminthes and the comparative efficiency test of nematicidal drugs in goats of Wallaga administrative region at Mechara settlement area. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- Bashir, A.L., Chishti, F.A., Hidayatullah, T.A., 2012. Survey of gastrointestinal helminthes parasites of slaughtered sheep and goats in Ganderbal, Kashmir. *Global Vet.*, 8(4), 338-341.
- Bayou, A., 1992. Prevalence of gastrointestinal helminthes of small ruminants in Buno province, Illubabor administrative region. DVM Thesis Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- Bersisa, K., Tigist, T., Teshale, S., Reta, D., Bedru, H., 2011. Helminths in sheep and goats in central Oromia during the dry season. *J. Anim. Vet. Adv.*, 10(14), 1845-1849.
- Bikila, E., Yeshitla, A., Worku, T., Teka, F., Benti, D., 2013. Epidemiology of gastrointestinal parasites of small ruminants in Gechi district, Southwest Ethiopia. *Adv. Biol. Res.*, 7(5), 169-174.
- Chanie, M., Begashaw, S., 2012. Assessment of the economic impact and prevalence of ovine Fasciolosis in Menz Lalo Midir district, Northeast Ethiopia. *Vet. World*, 5(5), 261-264.
- Colditz, I.G., Watson, D.L., Gray, G.D., Eady, S.J., 1996. Some relationships between age, immune responsiveness and resistance to the parasites in ruminants. *Int. J. Parasitol.*, 26, 869-877.

- CSA, 2012. Federal democratic republic of Ethiopia, central statistical agency agricultural sample survey. Report on livestock and livestock characteristics; Vol. II, Addis Ababa, March.
- Dagnachew, S., Asmare, A., Wudu, T., 2011. Epidemiology of gastrointestinal helminthiasis of small ruminants in selected sites of North Gondar zone, Northwest Ethiopia. *Ethiop. Vet. J.*, 15(2), 57-68.
- Dereje, G., 1992. Investigation of common gastrointestinal parasite of small ruminants in and around Wolaita Soddo. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- FAO, 2005. Ethiopia FAO's information system on water and agriculture. Rome, Italy.
- Fikru, R., Teshale, S., Reta, D., Yosef, K., 2006. Epidemiology of gastrointestinal parasites of ruminants in Western Oromia, Ethiopia. *Int. J. Appl. Res. Vet. Med.*, 1, 451-57.
- Fritsch, T., Kaufmann, J., Ptister, K., 1993. Parasite spectrum and seasonal epidemiology of gastrointestinal nematodes of small ruminants in Gambia. *Vet. Parasitol.*, 49(2-4), 271-283.
- Gamble, H.R., Zajak, A.M., 1992. Resistance of St. Croix lambs to *Haemonchus contortus* in experimentally and naturally acquired infections. *Vet. Parasitol.*, 41, 211-225.
- Gareth, W.H., 2009. Nematode parasites of small ruminants, camelids and cattle: Diagnosis with emphasis on anthelmintic efficacy and resistance testing. Australia and New Zealand Standard Diagnostic Procedures.
- Gebreyesus, M., 1986. Prevalence of gastrointestinal helminthes of small ruminants (sheep and goats) in Gonder administrative region. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre- Zeit, Ethiopia.
- Genene, R.A., 1994. Study of prevalence of ovine GIT helminthes in and around Kombolcha. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- Githigia, S.M., Thamsborg, S.M., Munyua, W.K., Maingi, N., 2001. Impact of gastrointestinal helminths on production in goats Kenya. *Small Rumin. Res.*, 42, 21-29.
- Hailelul, N., 2002. Study on prevalence of GIT helminthes of small ruminants in and around Wolayta Soddo, Southern Ethiopia. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia.
- Hansen, J., Perry, B., 1994. The epidemiology, diagnosis and control of helminth parasites of ruminants. A handbook. 2nd edition. ILRAD (International Laboratory for Research on Animal Diseases), Nairobi, Kenya.
- Keyyu, J.D., Kassuku, A.A., Kyvsgaard, N.C., Willingham, A.L., 2003. Gastrointestinal nematodes in indigenous zebu cattle under pastoral and nomadic management systems in the lower plain of Southern highlands of Tanzania. *Vet. Res. Comm.*, 27(5), 371-380.
- Kumba, F., Katjivena, H., Kauta, G., Lutaaya, E., 2003. Seasonal evolution of faecal egg output by gastrointestinal worms in goats on communal farms in Eastern Namibia. *Onderstepoort J. Vet. Res.*, 70(4), 265-271.
- Ntonifor, H., Shei, S., Ndaleh, N., Mbunkur, G., 2013. Epidemiological studies of gastrointestinal parasitic infections in ruminants in Jakiri, Bui division, Northwest region of Cameroon. *J. Vet. Med. Anim. Health*, 5(12), 344-352.
- Odoi, A., Gathuma, J.M., Gachui, C.K., Omore, A., 2007. Risk factors of gastrointestinal nematode parasite infections in small ruminants kept in smallholder mixed farms in Kenya. *BMC Vet. Res.*, 3, 6.
- Petros, A., Lakew, N., 2014. Prevalence of gastrointestinal parasites of small ruminants in Kuarit district, Northwest Ethiopia. *Afr. J. Basic Appl. Sci.*, 6(5), 125-130.
- Radiostits, O.M., Gray, C.C., Hincheliff, K.W., Constable, P.D., 2007. Veterinary medicine. A textbook of the disease of cattle, horses, sheep, pigs and goats. 10th edition. Saunders, Saunders Elsevier, London.
- Raza, M.A., Iqbal, Z., Jabbar, A., Yaseen, M., 2007. Point prevalence of gastrointestinal helminthiasis in ruminants in Southern Punjab. *Pakistan J. Helminthol.*, 81, 323-328.
- Sissay, M.M., 2007. Helminth parasites of sheep and goats in eastern Ethiopia: Epidemiology, and anthelmintic resistance and its management. Doctoral Thesis, Swedish University of Agricultural Sciences, Uppsala, Sweden.
- Soulsby, E.J.C., 1982. Helminthes, arthropods and protozoa of domestic animals. 7th edition, Lea and Faebiger, Philadelphia.
- Tefera, M., Batu, G., Bitew, M., 2011. Prevalence of gastrointestinal parasites of sheep and goats in and around Bedelle, Southwestern Ethiopia. *Internet J. Vet. Med.*, 8, 3-5.
- Teklye, B., 1991. Epidemiology of endoparasites of small ruminants in Sub-Saharan Africa. Proceedings of Fourth National Livestock Improvement Conference. Addis Ababa, Ethiopia; 13-15 November, 7-11.
- Temesgen, T., 2008. Study on prevalence of ovine gastrointestinal parasite in and around Bedelle. DVM Thesis, Haramaya University, Faculty of Veterinary Medicine, Haramaya, Ethiopia.

- Tesfaheywet, Z., 2012. Helminthosis of sheep and goats in and around Haramaya, Southeastern Ethiopia. J. Vet. Med. Anim. Health, 4(3), 48-55.
- Tesfaye, H., 1998. Ovine and bovine helminthiasis in Kelala (South Wollo). Ethiopian Vet Association Proceedings of the 12th Conference. Addis Ababa, Ethiopia; June, 30-34.
- UNECA, 2012. Report on livestock value chains in eastern and southern Africa: A regional perspective, Addis Ababa, Ethiopia, 19-21 November.
- Urquhart, G.M., Armour, J., Duncan, J.L., Dunn, A.M., Jennings, F.W., 1996. Veterinary parasitology. 2nd ed., Blackwell Science Ltd, UK, 141-205.
- Waruiru, R.M., Mutune, M.N., Otieno, R.O., 2005. Gastrointestinal parasite infections of sheep and goats in a semi-arid area of Machakos district, Kenya. Bull. Anim. Health Prod. Afr., 53(1), 25-34.

How to cite this article: Bedada, H., Gizaw, F., Negash, W., 2018. Preliminary study on small ruminant GIT helminthiasis in select arid and semi-arid pastoral and agro-pastoral areas of Afar region, Ethiopia. Scientific Journal of Veterinary Advances, 7(1), 210-217.

Submit your next manuscript to Sjournals Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in DOAJ, and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.sjournals.com

Sjournals
where the scientific revolution begins