

Contents lists available at [Sjournals](http://Sjournals.com)



Journal homepage: www.Sjournals.com



Short communication

Retrospective analysis of reproductive disease conditions among domestic ruminants in Sokoto, Nigeria

M.A. Umaru^{a,*}, A.A. Adeyeye^a, H.S. Garba^b, A. Bello^c, A. Abubakar^c

^aDepartment of Theriogenology and Animal Production, Usmanu Danfodiyo University, Sokoto, Nigeria.

^bDepartment of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Nigeri.

^cDepartment of Veterinary Anatomy, Usmanu Danfodiyo University, Sokoto, Nigeria.

*Corresponding author; Department of Theriogenology and Animal Production, Usmanu Danfodiyo University, Sokoto, Nigeria.

ARTICLE INFO

ABSTRACT

Article history:

Received 09 February 2013

Accepted 23 February 2013

Available online 28 February 2013

Keywords:

Livestock reproduction

Sheep

Goat

Cattle

Dystocia

Pregnancy toxaemia mastitis

A fifteen-year (1991 — 2005) study of reproductive cases in animals presented to the Usmanu Danfodiyo University Veterinary Teaching Hospital, Sokoto, were analyzed based on species, disease condition and sex using clinical case Files of Sheep, goat and cattle. Within the study period a total of 88 reproductive cases were handled out of which 53 (57.95 %) occurred in sheep, 32(36.36 %) goat and 5(5.68 %) cattle. Dystocia 23(26.13 %), Pregnancy toxaemia 11(12.50 %), mastitis 9(10.23 %), castration 5(5.68 %) and orchitis 3(3.41 %) were the diseases recorded. Reproductive cases were higher in females 77(87.5 %) than in mates 11(12.5 %). From the study, reproductive cases were most prevalent in sheep than in goats and cattle.

© 2013 Sjournals. All rights reserved.

1. Introduction

The production of livestock is dependant greatly on their well being particularly their reproductive performances. Disease conditions always impair livestock production (Akerejola *etal*, 1979; Lamorde, 1996). Apart from this, several other factors such as environment and nutrition, especially inadequate protein intake (Kumidiake *et al.*, 1981; Smith and Somade, 1994) decrease reproductive performance.

The knowledge of diseases prevalence give useful information on disease pattern and thus can be used in preventing diseases as well as formulating policies for future management of prevalent diseases. Although analysis of some common disease have been conducted in different parts of Nigeria (Esuruoso, 1972; Ugochukwu and Ephraim, 1985; Cadmus *et al.*, 2001), Ebbo *et al.* (2003) reported that there was little information on the prevalence of different livestock disease in Sokoto. However, only Wosu and Anene (1990) in Nsukka and Waziri *et al.* (2006) in Maiduguri have analyzed reproductive disease condition among ruminants.

This study is therefore aimed at determining the pattern of reproductive diseases encountered at the Usmanu Danfodiyo University Veterinary Teaching Hospital, Sokoto, within a fifteen-year period.

2. Materials and methods

The data obtained are from clinical case files of sheep, goat and cattle presented to the Usmanu Danfodiyo University Veterinary Teaching Hospital Sokoto, Nigeria from January 1991 - December 2005 (fifteen-year period). The data were analyzed based on species, disease condition and sex using percentage distribution in a tabular form.

3. Results

The analysis of various reproductive disease conditions in different sex is presented according to specie in Table I. A total of eighty-eight (88) reproductive cases were handled in domestic ruminants within the study period. Sheep had the highest prevalence of 51(57.95 %), followed by goats 32(36.36 %) then cattle 5(5.68 %). Dystocia 23(26.13 %) was the most prevalent reproductive condition, this is followed by Pregnancy toxemia 11(12.50 %), then Retained placenta 10(11.36 %), Mastitis 9(10.23 %); Parturient paresis 7(7.95%), Abortion 6(6.82 %); Castration 5(5.68 %); Vaginal prolapsed 4(4.55%); Stillbirth 4(4.55%); Orchitis 3(3.41 %); Uterine prolapse 3(3.41 %); Balanopostitis 2(2.27 %); and Phimosis 1(1.14 %). Reproductive cases in females 77(87.5 %) were more prevalent than males 11(12.5 %).

Table1

Disease condition	Cattle	Goat	Sheep	Total	Prevalence rate (%)
Dystocia	1	9	13	23	26.13
Preg. Toxaemia	-	1	10	11	12.50
Parturient Paresis	-	3	4	7	7.95
Retained placenta	1	4	5	10	11.36
Abortion	1	3	2	6	6.82
Stillbirth	-	1	3	4	4.55
Uterine	-	-	3	3	3.41
Vaginal Prolapse	-	1	3	4	4.55
Mastitis	1	3	5	9	10.23
Orchitis	-	2	1	3	3.41*
Balanopostitis	-	-1	2	2	2.27*
Castration	-	5	-	5	5.68*
Phimosis	1	-	-	1	1.14*
Total	5 (5.68%)	32(36.36%)	51(57.98)	88 (100)	100

* Prevalence of Male reproductive disease which sum-up to 12.5%.

4. Discussion

Higher prevalence of reproductive disease conditions in small ruminants (sheep and goat) over cattle has been recorded. The high number in the animal species of small ruminants is as a result of high number of cases handled in the hospital coupled with the low cost of rearing. This is consistent with the reports of Mohammed *et al.* (1995) and Waziri *et al.* (2006) but contradicts that of Wosu *et al.* (1990). The religious and socio-economic importance of sheep in Northern Nigeria may explain why most inhabitants of Sokoto keep them as it is used

during festive periods such as *eld-el kabir*, wedding and naming ceremonies. The number of cattle cases handled within the study period was low despite the report of Williams *et al.* (2000) that Sokoto state is the second largest cattle producing state in Nigeria. The low numbers of cattle cases handled may probably be due to high cost of rearing cattle in the metropolis as well as the high cost of transporting disease cattle from the rural parts of the state to the Veterinary Hospital Sokoto. Dystocia was the most prevalent reproductive condition. This is consistent with the reports of Wosu and Anene (1990) in Nsukka and Waziri *et al.* (2006) in Maiduguri. Dystocia is caused by twinning, poor feeding and management (Arthur *et al.* 1998). Free roaming of animals is a common scenario in Sokoto metropolis. Na-Allah *et al.* (2003) reported that most ruminant livestock kept by inhabitants of Sokoto are reared under semi-extensive system of production in which they, are allowed to roam freely; this probably explains the high occurrence of dystocia and pregnancy toxemia. Castration was the most frequent reproductive condition recorded in male goats. Castration is known to improve growth and thus meat quality in castrated ruminants.

Reproductive disease condition were more in females probably due to their unique position as essential reproductive vessels and the fact that females are reared for a longer periods than males. This was consistent with the report of Waziri *et al.* (2006) in Maiduguri.

5. Conclusion

The study showed that reproductive disease conditions were more frequent in sheep and female sexes were more vulnerable, with dystocia being the most prevalent reproductive condition encountered within the period.

References

- Akerejola, O.C., Schillhorn, V.T., Njoku, C.O., 1979. Ovine and Caprine diseases in Nigeria. *Bulletin of Animal Health and Production in Africa*, 27, 65 - 68.
- Arthur, G.H., Noakes, D.E., Persea, H., Parkinson, T.J., 1998. *Veterinary Reproduction and Obstetrics*, 7th Edition. W. B. Saunders Company, Philadelphia.
- Cadmus, S.J.B., Ogundipe, G.A., Abatan, A.A., Mwanla, P.G., Adesanya, G.A., Bulus, S.T., 2001. A study of clinical cases treated with economics of operation in a private Veterinary clinic in Abuja, Nigeria. *Tropical Veterinarian*, 19(1), 65 - 69.
- Ebbo, A.A., Agaie, B.M., Adamu, U., Daneji, A.I., Garba, H.S., 2003. Retrospective analysis of cases presented to the Usmanu Danfodio University Sokoto (1993 - 2002). *Nigerian Veterinary Journal* 24, 133 - 136.
- Esuruosu, G.O., 1972. Observation on an experimental veterinary clinic in Ikeja of Lagos. *Nigerian Veterinary Journal*, 1, 7 - 15.
- Kumidiaka, J., Osori, D.I.K., Ogwu, D., 1981. Incidence of genital abnormalities and physiological effect of genital pathology in indigenous cows. *Nigerian Veterinary Journal*, 9, 52 - 54.
- Lamorde, A.G., 1996. The role of veterinarians on a developing economy. *Nigerian Veterinary Journal* (special edition), 1(1), 106 - 111.
- Mohammed, A., Ahmed, B.C., 1995. An analysis of surgical cases at the University of Maiduguri Veterinary Teaching Hospital. *Annals of Borno*. 11/12, 303 - 308.
- Na-Allah, Y., Tukur, H.M., Maigandi, S.A., Daneji, A.I., 2003. Reproductive performance of Sheep and goat under traditional management system in Zamfara reserve north western Nigeria. *Sokoto Journal of Veterinary Science*, 5(1), 18 - 20.
- Smith, O.S., Somade, B., 1994. Interaction between nutrition and reproduction in farm animals. *IFS (International Foundation for science) Proceedings of a Regional Seminar on Animal Reproduction*. Jan 17 - 21. 1994.
- Ugochukwu, E.I., Ephraim, B.U., 1985. An analysis of prevalence of disease of domestic animals at the Veterinary Clinic Calabar Nigeria. *Nigerian Veterinary Journal*, 14(1), 42 - 44.
- Waziri, M.A., Adamu, A., Bukar, M.M., 2006. Analysis of reproductive cases handled at the State Veterinary Clinic, Maiduguri. Nigeria (1993 - 2005). *Nigerian Veterinary Journal*, 27(2), 54 - 59.
- Williams, A., Bzugu, P.M., Atsanda N.N., 2000. A retrospective study of diseases of ruminants at Maiduguri Nigeria. *Tropical Veterinarian*. 18, 23 - 28.
- Wosu, L.O., Anene, B.M., 1990. Incidence and seasonality of reproductive disease conditions in small ruminants in Nsukka area Nigeria. *Beitr Trop Landwirtsch Veterinarmed*, 28(5), 185 - 189.